

**This bibliography contains historic and current published scientific papers, contract reports, popular articles, posters and news items on topics of relevance to Tairāwhiti and to those participating in the Transition Advisory Group Meetings.**

**Prepared by Mike Marden  
October 2024.**

Many of the earlier published articles are not available in electronic form. For some of the later publications I have provided links to websites.

I've endeavoured to group these publications into themes. Some publications address more than one theme so are listed more than once.

Each publication will also contain a list of references some of which will be of relevance to this TAG.

This bibliography is by no means complete. Please feel free to suggest additional references of relevance.

Themes:

1. Previous reviews into impacts of erosion and floods on pastoral and forestry land pre-and post-Bola.
2. Costs of erosion, soil and carbon loss and flooding
3. NES-PF/CF
4. Historic mass movement and gully erosion
5. Vegetation & slope stability
6. Environmental effects of forestry including hydrology, sediment yields, water quality, & ecology
7. Alternative vegetative options/strategies for transitioning to a permanent forest cover
8. Forest management on highly erodible hill country
9. Slash management and riparian buffers
10. Changes in mean bed levels East Coast rivers
11. Sediment sources and delivery to stream channels
12. Social implications of erosion and forestry
13. Research posters
14. Popular articles
15. Newspaper articles/Radio/TV
16. Soils
17. Waiaapu Catchment, East Coast Region, North Island.
18. Waipaoa Catchment, East Coast Region, North Island
19. Waimata Catchment, East Coast Region, North Island

## **1. Previous reviews into impacts of erosion and floods on pastoral and forestry land pre-and post-Bola.**

National Water and Soil Conservation Organisation (NWASCO). (1970). *Wise Land Use and Community Development*. [Report of the technical committee of enquiry into the problems of the Poverty Bay-East Coast District of New Zealand. National Water and Soil Conservation Organisation.] Wellington, New Zealand: Water and Soil Division, Ministry of Works. 119p, also known as the Taylor Report, as this document sets the scene regarding land use, and in particular, erosion-related issues that have plagued this region since the late 1880—early 1920 through to present-day.

- Poverty Bay Catchment Board (1978). Report of land use planning and development study for erosion prone land in the East Cape Region. Section 1: The East coast (The Red Report).
- Water and Soil Directorate. (1987). *East Coast project review*. Wellington, New Zealand: Water and Soil Directorate, Ministry of Works, and Development. 123p & Appendices.
- Ministry of Agriculture and Fisheries (1988). Effects of erosion on the production and returns of East Coast Hill Country Farms. 19p.
- Office of Parliamentary Commissioner for The Environment. (1988). Inquiry into flood mitigation measures following Cyclone Bola. 59p.
- Ministry of Agriculture and Fisheries (1989). The effect of Cyclone Bola on hill country farms in the Gisborne -East coast Regions: Physical damage, Government assistance, cash flows and dept. 111p.
- East cape catchment Board (1989). East Coast Conservation Forestry Scheme: Review of Progress 1989. 159 % Appendices.
- Webber and Associates (1989). Cyclone Bola Agricultural Assistance Scheme: Economic and Sosial Impact Study. 86 p & Appendices.
- Ministry of Forestry. (1993). Report from Forestry Road Funding Task Force. 64p
- Ministry of Forestry. (1994). *A guide to the East Coast Forestry Project*. Wellington, New Zealand: Ministry of Forestry.
- Office of Parliamentary Commissioner for The Environment. (1994). *Sustainable Land Management and the East Coast Forestry Project*, 61 p. Wellington, New Zealand. <https://www.pce.parliament.nz/media/1546/sustainable-management-and-the-east-coast-forestry-project-dec-1994-small.pdf>.
- Ministry of Forestry. (1996). Forest Growing Investment on the East Coast. Ministry of Forestry Zone Studies.
- Ministry of Forestry. (1998). *The East Coast Forestry Project*. A discussion paper for the 1998 review. Wellington, New Zealand: Ministry of Forestry. 27p
- Bayfield, MA, Meister, AD (1998). The East Coast Forestry Project Review. Report to Ministry of Agriculture and Forestry. 52p.
- Ministry of Agriculture and Forestry. (2005). *The 2005 Review of the East Coast Forestry project: A Discussion Paper for the 2005 Review*. [MAF Discussion paper No. 37]. Wellington, New Zealand: Ministry of Agriculture and Forestry.
- Bayfield MA, Meister AD. (2005). *East Coast Forestry Project review*. Report to Ministry of Agriculture and Forestry, Wellington, NZ. 48p.
- Phillips, C.; & Marden, M. (2005). Reforestation schemes to manage regional landslide risk. Chapter 18 under Part III 'Management of Landslide Risk' in "Landslide Hazard and Risk". Edited by Glade, T.; Anderson, M.; Crozier, M. Publisher: J. Wiley & Son, p. 517-548. Includes summary of erosion control efforts before Cyclone Bola.
- Ministry for Primary Industries. (2017). *Erosion Control Funding Programme*.

<https://www.mpi.govt.nz/forestry/funding-tree-planting-research/closed-funding-programmes/erosion-control-funding-programme/>

## 2. Costs of erosion, soil and carbon loss and flooding

Basher L, McNeill S, Page M, Lynn I, Betts H, De Rose R, **Marden M**, Rosser B. (2013). Soil carbon stocks and changes: carbon losses from erosion. MPI Technical Paper no 2013/, 42 pp.

Krausse MK, Eastwood C, Alexander RR. (2001). Muddied waters: counting the national economic cost of soil erosion and sedimentation in New Zealand. Lincoln, NZ, Manaaki Whenua Press.

MfE (2008). Quantification of the flood and erosion reduction benefits, and costs, of climate change mitigation measures in New Zealand. Wellington, Ministry for the Environment.

Jones H, Clough P, Höck B, Phillips C. (2008). Economic costs of hill country erosion and benefits of mitigation in New Zealand: review and recommendation of approach. Scion Contract Report, prepared for the Ministry of Agriculture and Forestry.

Barry L, Upananda Paragahawewa U, Richard Yao R, Turner J. (2011). Valuing avoided soil erosion by considering private and public net benefits. NZARES Conference Tahuna Conference Centre, Nelson, New Zealand, 25-26 August 2011.

Dominati E, Mackay A. (2014). Using an ecosystem services approach to assess the cost of soil erosion. *Soil Horizons* 23.

Dominati EJ, Mackay A, Lynch B, Heath N, Millner I (2014). An ecosystem services approach to the quantification of shallow mass movement erosion and the value of soil conservation practices. *Ecosystem Services* 9: 204–215.

MPI (2015). Future requirements for soil management in New Zealand. Phases 1-3. MPI (2015). Ministry for Primary Industries Science Strategy Rautaki Putaiao.

**Krausse MK, Eastwood C, Alexander RR. (2001), Bayfield & Meister and Phillips and Marden (2005) also provide figures of the costs of remediation/repairs for Cyclone Bola.**

## 3. NES-PF

Ministry for Primary Industries (2017). National environmental standards for plantation forestry. <http://www.mpi.govt.nz/growing-and-harvesting/forestry/national-environmental-standards-for-plantation-forestry/> (accessed 16 February 2018)

Ministry for Environment (2017). About the National Environmental Standards for Plantation Forestry. <http://www.mfe.govt.nz/land/national-environmental-standards-plantation-forestry/about-standards> (accessed 16 February 2018)

Ministry for Primary Industries (2017). Plantation forestry erosion susceptibility classification: risk assessment for the National Environmental Standards for Plantation forestry. MPI Technical Paper No: 2017/47.

## 4. Historic mass movement and gully erosion

Basher L, Betts H, Lynn I, Marden M, McNeill S, Page M, Rosser B. 2013. The effect of earthflow erosion on soil carbon stocks. SLMACC project C09X1006 MPI Technical Paper No. 2012/ 55 pp.

- Betts, H.D.; DeRose, R.C. (1999): Digital elevation models as a tool for monitoring and measuring gully erosion. *Journal of Applied Earth Observation and Geoinformation*, 1(2): 91–101.
- Betts, H.D.; Trustrum, N.A.; DeRose, R.C. (2003): Geomorphic changes in a complex gully system measured from sequential digital elevation models, and implications for management. *Earth Surface Processes and Landforms* 28 (10): 1043-1058
- Bilderback, E. L., Pettinga, J. R., Litchfield, N. J., Quigley, M., Marden, M., Roering, J. J. and Palmer, A. S. 2015 Hillslope response to climate-modulated river incision in the Waipaoa catchment, East Coast North Island, New Zealand, *Bulletin of the Geological Society of America*, 127(1-2): 131-148.
- Blaschke, PM., Trustrum, NA., & Douglas L. Hicks (2000). Impacts of mass movement erosion on land productivity: a review. *Progress in Physical Geography* Vol. 24, No. 1, p21-52.
- Black, R. D. (1980): Upper Cretaceous and Tertiary geology of Mangatu State Forest, Raukumara Peninsula, New Zealand. *N. Z. Journal of Geology and Geophysics* 23: 293-312.
- Campbell, D. A. (1946): Down to the sea in slips. Soil Conservation and Rivers Control Council, Wellington, N. Z., Bulletin No. 5, 35p.
- Cerovski-Darriau, C., Roering, J.J., Marden, M., Palmer, A.S., Bilderback, E.L. 2014. Quantifying temporal variations in landslide-driven sediment production by reconstructing paleo landscapes using tephrochronology and lidar: Waipaoa River, New Zealand. *American geophysical Union, Geochemistry, Geophysics, Geosystems*, 15p. doi: 10.1002/2014GC005467.
- DeRose R, Basher L 2010. Measurement of riverbank and cliff erosion from sequential LIDAR and historical aerial photography. *Geomorphology* 126: 132–147.
- DeRose, R.C.; Gomez, B.; Marden, M.; Trustrum, N.A. (1998): Gully erosion in Mangatu Forest, New Zealand, estimated from digital elevation models. *Earth Surface Processes and Landforms* 23: 1045–1053.
- Dymond, J.R., and Hicks, D.L., 1986. Steepland erosion measured from historical aerial photographs. *Journal of Soil and Water Conservation*, v. 41, no. 4, p. 252–255.
- East Coast Project (1978): Report of land use planning and development study for erosion-prone land of the East Cape region, Section 1, The East Coast, May 1978. A report by the Poverty Bay Catchment Board, the 'Red Report'. 24p.
- Eyles, G.O. (1983): The distribution and severity of present soil erosion in New Zealand. *New Zealand Geographer* 39 (1): 12–28.
- Gage, M.; Black, R. D. (1979): Slope Stability and geological investigations at Mangatu State Forest. Technical paper No. 66. Forest Research Institute, New Zealand Forest Service. 37pp. and maps.
- Gibb, J. G. (1981): Coastal hazard mapping as a planning technique for Waiapu County, East Coast, North Island, New Zealand. He ripoata Whakature mo nga whenu papa-a-tai o te rohe o te kaunihera o Waiapu-Tairawhiti. Water & Soil technical publication.

Wellington. Ministry of Works & Development.

Gibbs, H. S. (1954): Soils and Agriculture of Matakaoa County. New Zealand. Soil Bureau – Bulletin 11. New Zealand department of scientific and industrial research, 52p.

Gomez, B.; Eden, D.N.; Hicks, D.M.; Trustrum, N.A.; Peacock, D.H.; Wilmhurst, J. (1999): Contribution of floodplain sequestration to the sediment budget of the Waipaoa river, New Zealand. In: Marriott, S.B., Alexander, J. (eds) (1999): Floodplains: Interdisciplinary approaches. Special Publications, 163. London, Geological Society of London. Pp. 69–88.

Griffiths, G.A. (1982): Spatial and temporal variability in suspended sediment yields of North Island basins, New Zealand. *Water Resources Bulletin* 8 (4): 575-583.

Harmsworth, G.; Warmenhoven, T.; Pohatu, P. (2001): A Waiapu catchment management strategy: To stimulate interest and planning in the Waiapu, a strategy to underpin environmental management, economic development and sustain cultural values. A document for community and stakeholder discussion. Te Whare Wananga o Ngati Porou and Manaaki Whenua. 11p.

Harmsworth, G.; Warmenhoven T.; Pohatu, P.; Page, M. (2002): Waiapu catchment Technical Report: Maori community goals for enhancing ecosystem health. Foundation for Research, Science, and Technology (FRST) contract TWWX0001. Landcare Research report LC 0102/100 for Te Whare Wananga o Ngati Porou, Ruatorea (unpublished). 185p.

Hicks, D. M.; Griffiths, G. A. (1992): Sediment load. In: Mosley, M. P. (ed.): Waters of New Zealand. Wellington, New Zealand. Hydrological Society. Pp. 229–248.

Hicks, D.M.; Gomez, B.; Trustrum, N.A. (2000): Erosion thresholds and suspended sediment yields, Waipaoa River Basin, New Zealand. *Water Resources Research* 36 (4), 1129–1142.

Hicks, M.; Shankar, U.; McKerchar, A. (2003): Sediment yield estimates: a GIS tool. *Water & Atmosphere* 11(4): 26-27.  
<http://www.niwascience.co.nz/pubs/wa/ma/11-4/estimates>

Hughes, S.; Hughes, I. (1990): The Waiapu Mountains of the Raukumara Range. Gisborne, Eastland Promotion Council/Te Rau Press.

Kasai, M.; Marutani, T.; Reid L.M.; Trustrum, N.A. (2001): Estimation of temporally averaged sediment delivery ratio using aggradational terraces in headwater catchments of the Waipaoa River, North Island, New Zealand. *Earth Surface Processes and Landforms* 26, 1–16.

Kasai, M.; Brierley, G. J.; Page, M. J.; Marutani, T.; Trustrum, N. A. (2005): Impacts of land use change on patterns of sediment flux in Weraamaia catchment, New Zealand. *Catena* 64 (1): 27-60.

Kasai, M. (2006): Channel processes following land use changes in a degrading steep headwater stream in North Island, New Zealand. *Geomorphology* 81 (3-4): 421-439.

Kelliher, F.M.; Marden, M.; Watson, A.J.; Arulchelvam, I.M. (1995). Estimating the risk of landsliding using historical extreme flood data (Note). *Journal of Hydrology* 33: 123-129.

- Kenny, J. A. (1980): Geology of the Ihungia catchment, Raukumara Peninsula. M Sc, Dept. of Geology, University of Auckland.
- Kniskern, T.A.; Kuehl, S. A. (2004): Sediment dispersal and deposition on the Waiapu River Shelf, N.Z., implications for sediment transport mechanisms and event preservation. American Geophysical Union Meeting, San Francisco, CA, December 2004.
- Kniskern, T. A.; Harris, C. K.; Kuehl, S. A. (2006): Sediment deposition on the Waiapu River Shelf, N.Z., implications for sediment transport mechanisms and event Preservation. Ocean Sciences Meeting, Honolulu, Hawaii.
- Kuehl, S.A.; Pratson, L.; Addington, L.; Gerald, L.; Gerber, T.; Kniskern, T.; Miller, A.; Liu, P.; Carter., L.; Orpin, A. (2006): Contrasting Shelf Sediment Dispersal off Small Mountainous rivers: The Waipaoa and Waiapu Rivers, NZ. Ocean Sciences Meeting, Honolulu, Hawaii.
- Landcare Research (2001): Waiapu project GIS tables. Palmerston North, Landcare Research.
- Leathwick, J. R.; Clarkson, B. D.; Burns, B. R. Innes, J. G. Smale, M. C. (1995): Waiapu Ecological District. Survey report for the Protected Natural Areas Programme. NZ PNA Programme No. 31. Department of Conservation, Gisborne.177 p.
- Liébault, F.; Gomez, B.; Page, M.J.; Marden, M.; Peacock, D.H.; Richard, D.; Trotter, C.M. (2005): Land-use change, sediment production and channel response in upland regions. *River Research and Applications* 21, 739–756.
- Marden, M. (2011). Sedimentation history of Waipaoa Catchment, Landcare Research, Gisborne.
- Marden, M.; Arnold, G.; Gomez, B.; Rowan, D. (2005). Pre-and post-reforestation gully development in Mangatu Forest, East Coast, North Island, New Zealand. *River Research and Application Special Issue* 21, 757-771.
- Marden M, Arnold G, Seymour A, Hambling R. (2012). History and distribution of steepland gullies in response to land use change, East Coast Region, North Island, New Zealand. *Geomorphology*, 153: 81-90. <https://doi.org/10.1016/j.geomorph.2012.02.011>
- Marden M, Betts H, Arnold G, Hambling R. (2008). Gully erosion and sediment load: Waipaoa, Waiapu and Uawa rivers, eastern North Island, New Zealand. In: Schmidt J, Cochrane T, Phillips C, Elliott S, Davies T, Basher L eds Sediment dynamics in changing environments. International Association of Hydrological Sciences Publication 325. Wallingford, Oxfordshire, UK, IAHS. Pp. 339–350.
- Marden, M; Fuller, I; Herzig, A; Betts, H. (2018). Badass gullies: Fluvio-mass-movement gully complexes in New Zealand’s East Coast region, and potential for remediation. *Geomorphology* 307, 12-23. <https://doi.org/10.1016/j.geomorph.2017.11.012>
- Marden M, Herzig A, Arnold G. (2011). Gully degradation, stabilisation and effectiveness of reforestation in reducing gully-derived sediment, East Coast region, North Island, New Zealand. *Journal of Hydrology (New Zealand)*, 50 (1), 19-36.

- Marden, M., Herzig, A. and Basher, L. 2014 Erosion process contribution to sediment yield before and after the establishment of exotic forest: Waipaoa catchment, New Zealand, *Geomorphology*, 226: 162-174.
- Marden, M.; Phillips, C.J.; Rowan, D. 1991: Declining soil loss with increasing age of plantation forests in the Uawa Catchment, East Coast Region, North Island. In proceedings of International Conference on Sustainable Land Management, Nov. 17-23, Napier, New Zealand. pp. 358-361.
- Marden, M.; Rowan, D.; Phillips, C.J. (1995). Impact of cyclone-induced landsliding on plantation forests and farmland in the East Coast Region: A lesson in risk management? *In Proceedings XX IUFRO World Congress, Finland, Technical Session on Natural Disasters in Mountainous Areas.* pp. 133-145.
- Marden M, Rowan D, Phillips C. (2008). Recurrent displacement of a forested earthflow and implications for forest management, East Coast Region, New Zealand. In: Schmidt J, Cochran T, Phillips C, Elliott S, Davies T, Basher L eds *Sediment dynamics in changing environments.* International Association of Hydrological Sciences Publication 325. Wallingford, Oxfordshire, UK, IAHS. Pp. 491–501.
- Marden, M., & Seymour, A. (2022). Effectiveness of vegetation mitigation strategies in the restoration of fluvial and fluvio-mass movement gully complexes over 60-years, East coast region, North Island, *New Zealand Journal of Forestry Science* 52:19. <https://doi.org/10.33494/nzifs522022x226x>
- Page, M., Trustrum, N & Gomez, B. (2000). Implications of a century of Anthropogenic erosion for future land use in the Gisborne-East Coast region of New Zealand. *New Zealand Geographer* Vol. 56, No. 2, p9-20.
- Page, M.J.; Reid, L.M.; Lynn, I.H. 1999: Quantifying sediment production from Cyclone Bola landslides, Waipaoa catchment. *New Zealand Journal of Hydrology* 38: 289-308. <https://www.jstor.org/stable/43944823>
- Parkner T, Page M, Marden M, Marutani T 2007. Gully systems under undisturbed indigenous forest, East Coast region, New Zealand. *Geomorphology* 84(3–4): 241–253.
- Reid, L.M., & Trustrum, N.A. (In Press). Sediment budgets and land management planning: examples from NZ. *Journal of Environmental Planning and Management*.
- Reid LM, Page MJ (2003). Magnitude and frequency of landsliding in a large New Zealand catchment. *Geomorphology* 49(1–2): 71–88.
- Phillips, C.J. (1988): Geomorphic effects of two storms on the upper Waitahaia River catchment, Raukumara Peninsula, New Zealand. *Journal of Hydrology (NZ)* 27 (2): 99-112.
- Phillips, C.J. (1989). Geomorphic effects of Cyclone Bola 1988 - A note. *Journal of Hydrology (NZ)* 28 (2): 142-146.



Trotter C. (1988). Cyclone Bola: the inevitable disaster. *New Zealand Engineering* 43(6): 13–16.

Trustrum, N.A., Gomez, B., Page, M.J., Reid, L.M., & Hicks, D.M. (1999). Sediment production, storage, and output: The relative role of large magnitude events in steepland catchments. *Zeitschrift für Geomorphologie*, Suppl. Bd. 115, p71-86.

## 5. Vegetation & slope stability

Allsop, F. 1973 *The Story of Mangatu*, Government Printer, Wellington.

Bergin, D.O.; Kimberley, M.O.; Marden, M. 1993: How soon does regenerating scrub control erosion? *New Zealand Forestry*, August 1993. Pp. 38–40.

Bergin, D.O.; Kimberley, M.O.; Marden, M. 1995: Protective value of regenerating tea-tree stands on erosion-prone hill country, East Coast, North Island, New Zealand. *New Zealand Journal of Forestry Science*: 3–19.

Ekanayake, J.C.; Marden, M.; Watson, A.; Rowan, D. (1997). Tree roots and slope stability: A comparison between *Pinus radiata* and kanuka. *New Zealand Journal of Forestry Science* 27(2): 216-233.

Ekanayake, J.C.; Phillips, C.J.; Marden, M. (1999). A comparison of methods for stability analysis of vegetated slopes. *In* proceedings of the First Asia-Pacific Conference on Ground and Water Bioengineering for Erosion Control and Slope Stabilisation. April 19-21 Manila, The Philippines p.441-418.

Gage M, Black RD (1979) 'Slope-stability and geological investigations at Mangatu state forest.' Forest Research Institute, No. 66.

Glade T (2003) Landslide occurrence as a response to land use change: a review of evidence from New Zealand. *Catena* 51, 297-314.

Herzig A, Dymond J, Marden M. (2011). A gully-complex model for assessing gully stabilisation strategies. *Geomorphology*. <http://doi:10.1016/j.geomorph.2011.06.012>

Hicks DL (1991) Erosion under pasture, pine plantations, scrub, and indigenous forest: a comparison from cyclone Bola. *New Zealand Forestry* 1991, 21-22.

Kelliher, F.M.; Marden, M.; Watson, A.J.; Arulchelvam, I.M. (1995). Estimating the risk of landsliding using historical extreme river flood data. *Journal of Hydrology (New Zealand)* V.33 (2): 123-129.

Marden, M.; Rowan, D. (1993): Protective role of vegetation on Tertiary terrain before and during Cyclone Bola, East Coast, North Island, New Zealand. *New Zealand Journal of Forestry Science* 23(3): 255-263.  
[https://www.scionresearch.com/\\_data/assets/pdf\\_file/0011/59753/NZJFS2331993MARDON255\\_263.pdf](https://www.scionresearch.com/_data/assets/pdf_file/0011/59753/NZJFS2331993MARDON255_263.pdf)

Marden, M. (1994). Value of environmental monitoring of forest management under the Resource Management Act: Mangatu Forest. *New Zealand Forestry*, May, 39-41.



- Marden, M. (1998). New focus favoured for gully erosion. *Gisborne Herald*, 6 January 1998.
- Marden, M. (1998). Area's erosion may shock world. Land use policies counteractive. *Gisborne Herald*, Wednesday, 30 December 1998.
- Marden M (2004). Future-proofing erosion-prone hill country against soil degradation and loss during large storm events: have past lessons been heeded? *New Zealand Journal of Forestry* 49, 11-16.
- Marden, M, Basher, L, Phillips, C. (2015). Should detailed terrain stability or erosion susceptibility mapping be mandatory in erodible steeplands? *New Zealand Journal of Forestry* 59, 4, 32-42.
- Marden, M; Rowan, D; Lambie, S. (2016). Root development and whole-tree allometry of juvenile trees of five seed-lots of *Pinus radiata* D. Don: implications for forest establishment on erosion-prone terrain, East Coast region, North Island, New Zealand. *New Zealand Journal of Forestry Science* 46:24. <https://doi.org/10.1186/s40490-016-0082-y>
- Marden, M., Rowan, D., Watson, A. (2023). Effect of changes in forest water balance and inferred root reinforcement on landslide occurrence and sediment generation following *Pinus radiata* harvest on Tertiary terrain, eastern North Island, New Zealand. *New Zealand Journal of Forestry Science* 53:4. <https://doi.org/10.33494/nzjfs532023x216x>
- Marden M. (2012). Effectiveness of reforestation in erosion mitigation and implications for future sediment yields, East Coast catchments, New Zealand: A review. *New Zealand Geographer*, 68(1): 24-35. <https://doi.org/10.1111/j.1745-7939.2012.01218.x>
- Marden M, Phillips C 2011. Poplar and willow growth during their formative years: preliminary findings from new field trials. *EnviroLink Advice Grant: 907-GSDC83*, 23 pp.
- Murton BJ 1968. Mapping the immediate pre-European vegetation on the east coast of the North Island of New Zealand. *The Professional Geographer* 20: 262–264.
- Phillips C, Marden M, Basher L. (2012). Plantation forest harvesting and landscape response - what we know and what we need to know. *New Zealand Journal of Forestry*, 56(4): 4-12. [http://www.nzjf.org/contents.php?volume\\_issue=j56\\_4](http://www.nzjf.org/contents.php?volume_issue=j56_4)
- O'Loughlin, C.L.; Zhang, X.B. 1986: The influence of fast-growing conifer plantations on shallow landsliding and earthflow movement in New Zealand's steeplands. *In Proceedings of the 18<sup>th</sup> IUFRO World Congress, Ljubljana, Yugoslavia, September, Division 1, Volume 1, pp. 217-226.*
- O'Loughlin, C. (1995). The sustainable paradox – an examination of the plantation effect – a review of the environmental effects of plantation forestry in New Zealand. *New Zealand Forestry*, 39(4), 3-8.
- O'Loughlin, C., Watson, A. (1979). Root-wood strength deterioration in radiata pine after clearfelling. *New Zealand Journal of Forestry Science*, 9(3), 284-293. [https://www.scionresearch.com/data/assets/pdf\\_file/0006/59136/NZJFS931979OLOUGHLIN284\\_293.pdf](https://www.scionresearch.com/data/assets/pdf_file/0006/59136/NZJFS931979OLOUGHLIN284_293.pdf).

- O'Loughlin, C.L. 1984: Effectiveness of introduced forest vegetation for protection against landslides and erosion in New Zealand steeplands. Symposium on Effects of Forest Land Use on Erosion & Slope Stability, Hawaii
- O'Loughlin CL. (1974). The effect of timber removal on the stability of forest soils. *Journal of Hydrology (NZ)* 13:121–34.
- Pearce AJ, O'Loughlin CL, Jackson RJ, Zhang XB 1987. Reforestation: on-site effects on hydrology and erosion, eastern Raukumara Range, New Zealand. In: *Forest Hydrology and Watershed Management, International Association of Hydrological Sciences Publication 167*. Vancouver, IAHS. Pp. 489–497.
- Phillips, C. J. 1988 Geomorphic effects of two storms on the Upper Waitahaia River catchment, Raukumara Peninsula, New Zealand, *Journal of Hydrology*, 27(2): 1988.
- Phillips, C. J. 1989 Geomorphic effects of Cyclone Bola 1988: A note, *Journal of Hydrology*, 28(2): 142-146.
- Phillips, C; Marden, M; Rowan, D. (1989). Planning for Forestry after Cyclone Bola-a comment. *New Zealand Forestry*, November 16-17.
- Phillips CJ, Marden M, Pearce AJ 1991. Effectiveness of reforestation in prevention and control of landsliding during large cyclonic storms. In: *Proceedings 19<sup>th</sup> International Union of Forestry Research Organisations*, Montreal. Pp. 358–361.
- Phillips C, Marden M 2006. Reforestation schemes to manage regional landslide risk. In: Glade T, Anderson MG, Crozier MJ eds *Landslide hazard and risk*. Chichester, UK, John Wiley and Sons. Pp. 731–752.
- Preston NJ, Crozier MJ 1999. Resistance to shallow landslide failure through root-derived cohesion in East Coast hill country soils, North Island, New Zealand. *Earth Surface Processes and Landforms* 24: 665–675.
- Phillips, C.J.; Watson, A.J. (1994). Structural tree root research in New Zealand. Manaaki Whenua Press, Landcare Research Science Series No. 7., Lincoln, New Zealand. 71 pp.
- Phillips, C.; Marden, M.; Miller, D. (2000). Review of plant performance for erosion control in the East Coast Region. LCR Contract Report LC9900/111 (unpublished) 128p.
- Poole AL. (1960). Protection forests in New Zealand, and a Poverty Bay example. *New Zealand Geographer* 16: 115–130.
- Quilter SJ, Korte CJ, Smith DR 1993. Low cost revegetation of slips near Gisborne. *Proceedings of the New Zealand Grassland Association* 55: 187–191.
- Rhodes, D. 2001 Rehabilitation of deforested steep slopes on the East Coast of New Zealand's North Island, <http://www.fao.org/docrep/004/y2795e/y2795e06.htm>, Website accessed 01/08/2014.

Smale MC, McLeod M, Smale PN 1997. Vegetation and soil recovery on shallow landslide scars in tertiary hill country, East Cape Region, New Zealand. *New Zealand Journal of Ecology* 21: 31–41.

Thompson RC, Luckman PG (1993) Performance of biological erosion control in New Zealand soft rock hill terrain. *Agroforestry Systems* **21**, 191-211.

Watson, A.J.; Marden, M.; Rowan, D. (1995). Tree species performance and slope stability. *In* *Vegetation and Slopes*, Ed. D.H. Barker. pp.161-171. Thomas Telford Press, London. <https://doi.org/10.1680/vasspae.20313.0018>

Watson, A.J.; Phillips, C.J.; Marden, M. (1999). Root strength, growth, and rates of decay: root reinforcement changes of two tree species and their contribution to slope stability. *Plant and Soil (00)*: 1-9. <https://doi.org/10.1186/s40490-016-0082-y>

Watson, A. J.; O'loughlin, C.L. (1985). Morphology, strength and biomass of manuka roots and their influence on slope stability. *New Zealand Journal of Forestry Science* 15: 337–348.

Watson, A.J.; Marden, M; Rowan, D. (1997). Root-wood strength deterioration in kanuka after clear felling. *New Zealand Journal of Forestry Science* 27: 205–215.

## **6. Environmental effects including hydrology, sediment yields, water quality, & ecology**

Baillie, B.R. and Neary, D.G. (2015). Water Quality in New Zealand's Planted Forest: A Review. *New Zealand Journal of Forestry Science*, 45(7). doi:10.1186/s40490-015-0040-0.

Davies-Colley, R.; Scarsbrook, M.; Marden, M. (1999). What lives in muddy east coast streams? Research in soft-rock catchments. *Water & Atmosphere* 7 (3): 7.

Dons, A. (1987). Hydrology and sediment regime of a pasture, native forest, and pine forest catchment in the central North Island, New Zealand. *New Zealand Journal of Forestry Science*, 17: 2-3, 161-178.

Duncan, M.J. (1995). Hydrological impacts of converting pasture and gorse to pine plantation, and forest harvesting, Nelson, New Zealand. *Journal of Hydrology (New Zealand)* 34: 15-41.

Fahey, B.D. (1964). Throughfall and interception of rainfall in a stand of radiata pine. *Journal of Hydrology (New Zealand)* 3: 17-26.

Fahey BD, Marden M (2000). Sediment yields from a forested and a pasture catchment, coastal Hawkes Bay, North Island, New Zealand. *Journal of Hydrology (NZ)* 39, 49-63.

Gomez, B.; Eden, D.N.; Hicks, D.M.; Trustrum, N.A.; Peacock, D.H.; Wilmhurst, J. 1999: Contribution of floodplain sequestration to the sediment budget of the Waipaoa river, New Zealand. *In*: Marriott, S.B., Alexander, J. eds *Floodplains: Interdisciplinary approaches. Special Publications, 163*. London, Geological Society of London. Pp. 69–88.

- Hicks, D.M. (1988). Differences in suspended sediment yield from basins established in pasture and in exotic forest. In: Proceedings of NZ Hydrological Society Symposium, Dunedin, August 1988 5p.
- Hicks, D.M.; Gomez, B.; Trustrum, N.A. 2000: Erosion thresholds and suspended sediment yields, Waipaoa River basin, New Zealand. *Water Resources Research* 36 (4): 1129–1142.
- Kelliher, F.M.; Whitehead, D.; Pollock, D.S. (1992). Rainfall interception by trees and slash in a young *Pinus radiata* D.Don stand. *Journal of Hydrology* 131:187-204.  
[https://doi.org/10.1016/0022-1694\(92\)90217-J](https://doi.org/10.1016/0022-1694(92)90217-J)
- MacLaren, J.P. (1996). Environmental effects of planted forests. FRI bulletin No. 98.
- Marden, M. 2011. Sedimentation history of Waipaoa Catchment; Envirolink project 1015-GSDC96. 48p.
- O’Loughlin, C.L. (1977). Plantation forestry: hydrological and erosion aspects. *New Zealand Journal of Forestry*, 22(2): 238-241.
- O’Loughlin, C.L. (1994). The forest and water quality relationship. *New Zealand Forestry*, 39: 3, 26-30.
- Parkyn SM, Davies-Colley R, Scarsbrook MR, Halliday NJ, Nagels JW, Marden M, Rowan D. (2006). Pine afforestation and stream health: a comparison of land-use in two soft rock catchments East cape New Zealand. *New Zealand Natural Sciences* 31: 113–135.
- Pearce, A.J.; O’Loughlin, C.L.; Jackson, R.J.; Zhang, X.B. (1987). Reforestation: on-site effects on hydrology and erosion, eastern Raukumara Range, *New Zealand. International Association of Hydrological Sciences Publication* 167: 489-497.
- Quinn JM, Stroud M J (2002). Water quality and sediment and nutrient export from New Zealand hill-land catchments of contrasting land use. *New Zealand Journal of Marine and Freshwater Research* 36, 409-429.
- Reid, D.J., Quinn, J.M. and Wright-Stow, A.E. (2010). Responses of Stream Macroinvertebrate Communities to Progressive Forest Harvesting: Influences of Harvest Intensity, Stream Size and Riparian Buffers. *Forest Ecology and Management*, 260: 1804–1815
- Richardson, J.; Jowett, I. G. (2002): Effects of sediment on fish communities in East Cape streams, North Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* (36): 431-442. <http://www.rsnz.org/publish/nzjmfr/2002/033.php>
- Rowe LK, Marden M, Rowan D 1999. Interception and throughfall in a regenerating stand of kanuka (*Kunzea ericoides* var. *ericoides*), East Coast region, North Island, New Zealand, and implications for soil conservation. *Journal of Hydrology (NZ)* 38(1): 29–48.
- Warmenhoven, T. (2000). “River research reveals serious sediment pollution”. Te Runanga O Ngati Porou newsletter, NatiLink, October 2000.

Whitehead, D.; Kelliher, F.M. (1991). A Canopy water balance model for a *Pinus radiata* stand before and after thinning. *Agricultural and Forest Meteorology* 55: 109-126.

## 7. Alternative species options/strategies for transitioning to a permanent forest cover

- Bergin, D.O.; Kimberley, M.O.; Marden, M. (1995). Protective value of regenerating tea tree stands on erosion-prone hill country, East Coast, North Island, New Zealand. *New Zealand Journal of Forestry Science* 25(1): 3-19.
- Bergin, D. O; Kimberley, M.O.; Marden, M. (1993). How soon does regenerating scrub control erosion? *New Zealand Forestry*, August 1993.
- Lambie, SM., Awatere, S., Daigneault, A., Kirschbaum, MUF., **Marden, M.**, Soliman, T., Spiekermann, RI., Walsh, PJ. (2021). Trade-offs between environmental and economic factors in conversion from exotic pine production to natural regeneration on erosion prone land. *New Zealand Journal of Forestry Science*. 51:14. <https://doi.org/10.33494/nzjfs512021x163x>
- Lambie, S., & **Marden, M.**, Kirschbaum, M., Soliman, T., Walsh, P. (2018). Best options for land use following radiata harvest in the Gisborne District under climate change: Literature review SLMACC 405415 MPI Technical Paper No: 2018/46. LC3190. 57p. ISBN No: 978-1-77665-952-4 (online) ISSN No: 2253-3923 (online).
- Marden, M., Rowan, D. and Phillips, C. (2005). ‘Stabilising characteristics of New Zealand indigenous riparian colonising plants. *Plant and Soil*, 278, 95-105. <https://doi.org/10.1007/s11104-004-7598-2>
- Marden M, Lambie S. (2015) Plot-based, growth performance of space-planted manuka (*Leptospermum scoparium*) on marginal land, and vulnerability to erosion. Ministry of Primary Industries Technical Paper No. 2015/19. 35p. ISBN No: 978-0-908334-79-7 (online).
- Marden, M., Lambie, S., & Phillips, C. (2020). Potential effectiveness of low-density plantings of mānuka (*Leptospermum scoparium*) as an erosion mitigation strategy in steeplands, northern Hawke’s Bay, New Zealand. *New Zealand Journal of Forestry Science*, 50:10. <https://doi.org/10.33494/nzjfs502020x82x>
- Marden, M., Lambie, S., & Rowan, D. (2018a). Root system attributes of 12 juvenile indigenous early-colonising shrub and tree species with potential for mitigating erosion in New Zealand. *New Zealand Journal of Forestry Science*, 48:11. <https://doi.org/10.1186/s40490-018-0115-9>
- Marden, M; Lambie, S; Phillips, C. (2018b). Biomass and root attributes of 8 of New Zealand’s most common indigenous evergreen conifer and broadleaved forest species during the first 5 years of establishment. *New Zealand Journal of Forestry Science* 48:9. <https://doi.org/10.1186/s40490-018-0113-y>.
- Marden, M., Rowan, D., Watson, A. (2023). Effect of changes in forest water balance and inferred root reinforcement on landslide occurrence and sediment generation following *Pinus radiata* harvest on Tertiary terrain, eastern North Island, New Zealand. *New Zealand Journal of Forestry Science* 53:4. <https://doi.org/10.33494/nzjfs532023x216x>

- Phillips, C.; & **Marden, M.** (2005). Reforestation schemes to manage regional landslide risk. Chapter 18 under Part III ‘Management of Landslide Risk’ in “Landslide Hazard and Risk”. Edited by Glade, T.; Anderson, M.; Crozier, M. Publisher: J. Wiley & Son, p. 517-548.
- Phillips, C.J., Marden, M., Lambie, S.M., Watson, A., Ross, C., Fraser, S. (2012). Observations of below-ground characteristics of young redwood trees (*Sequoia sempervirens*) from two sites in New Zealand – implications for erosion control. *Plant and Soil*, 1-16. <http://dx.doi.org/10.1007/s11104-012-1286-4>
- Spiekermann, R & **Marden, M.** (2018). Best options for land use following radiata harvest in the Gisborne District under climate change: Spatial analysis of erosion susceptibility in plantation forests, East Coast Region SLMACC 405415. MPI Technical Paper No: 2018/47. ISBN No: 978-1-77665-953-1 (online) ISSN No: 2253-3923 (online).
- Trotter, C., Tate, K., Scott, N., Townsend, J., Wilde, H., Lambie, S., Marden, M., Pinkney, T. (2005). Afforestation/reforestation of New Zealand marginal pasture lands by indigenous shrublands: the potential for Kyoto Forest sinks. *Annals of Forest Science* 62, 865-871.

## 8. Forest management on highly erodible hill country

- Amishev, D, Basher L, Phillips C, Hill S, Marden M, Bloomberg M, Moore J. (2013). New Forest Management Approaches to Steep Hills. Prepared for the Ministry for Primary Industries, 109 pp. ISBN No: 978-0-478-43786-7 (online) ISSN No: 2253-3923 (online).
- Basher, L., Harrison, D., Phillips, C., Marden, M. (2015). What do we need for a risk management approach to steepland plantation forests in erodible terrain? *NZ Journal of Forestry*, Vol. 60, No. 2
- Bloomberg, M. (2015). Erosion susceptibility classification and analysis of erosion risks for plantation forestry – response to Marden et al. *NZ Journal of Forestry*, Vol. 60, No. 2
- Funk JM, Field CB, Kerr S, Daigneault, A 2014. Modelling the impact of carbon farming on land use in a New Zealand landscape. *Environmental Science & Policy* 37: 1–10.
- Lambie, S.M., Awatere, S., Daigneault, A., Kirschbaum, M.U.F., Marden, M., Soliman, T., Spiekermann, R.I., Walsh, P.J. (2021). Trade-offs between environmental and economic factors in conversion from exotic pine production to natural regeneration on erosion prone land. *New Zealand Journal of Forestry Science*. 51:14. <https://doi.org/10.33494/nzifs512021x163x>
- Marden, M. (1994). Value of environmental monitoring of forest management under the Resource Management Act: Mangatu Forest. *New Zealand Forestry*, May, 39-41.
- Marden M (2004) Future-proofing erosion-prone hill country against soil degradation and loss during large storm events: have past lessons been heeded? *New Zealand Journal of Forestry* 49, 11-16.
- Marden, M. (2003). Gully erosion, the cancer of Waiapu catchment: 901 reasons to act now. *Tairāwhiti Conservation Quorum*, Issue 32, Winter 2003.
- Marden M. (2012). Effectiveness of reforestation in erosion mitigation and implications for future sediment yields, East Coast catchments, New Zealand: A review. *New Zealand Geographer*, 68(1): 24-35. <https://doi.org/10.1111/j.1745-7939.2012.01218.x>



- Marden, M., Basher, L., Phillips, C. (2015). Should detailed terrain stability or erosion susceptibility mapping be mandatory in erodible steeplands? *New Zealand Journal of Forestry* 59, 4, 32-42.
- Marden, M., & Seymour, A. (2022). Effectiveness of vegetation mitigation strategies in the restoration of fluvial and fluvio-mass movement gully complexes over 60-years, East coast region, North Island, *New Zealand*. *New Zealand Journal of Forestry Science* 52:19. <https://doi.org/10.33494/nzjfs522022x226x>
- Payn, T., Phillips, C., Basher, L., Baillie, B., Garrett, L., Harrison, D., Heaphy, M., and Marden, M. (2015). Improving management of post-harvest risks in steepland plantations. *NZ Journal of Forestry*, Vol. 60, No. 2.
- Phillips C, Marden M, Basher L. (2012). Plantation forest harvesting and landscape response - what we know and what we need to know. *New Zealand Journal of Forestry*, 56(4): 4-12. [http://www.nzjf.org/contents.php?volume\\_issue=j56\\_4](http://www.nzjf.org/contents.php?volume_issue=j56_4)
- Phillips, C; Marden, M; Rowan, D. (1989). Planning for Forestry after Cyclone Bola-a comment. *New Zealand Forestry*, November 16-17.
- Phillips, C., Marden, M., Basher, L.R. (2017). Geomorphology and forest management in New Zealand's erodible steeplands: an overview. *Geomorphology*, 307, 109-121. <https://doi.org/10.1016/j.geomorph.2017.07.031>
- Phillips, C; Basher, L; Marden, M. (2017). A risk matrix for storm-initiated forestry-related landslides and debris flows in the Gisborne region. Landcare Research Contract Report LC2711 Prepared for Gisborne District Council.30p.
- Phillips, C; Basher, L; Marden, M. (2016). Research and monitoring advice on environmental impacts of forestry in the Gisborne-East Coast region. Landcare Research Report LC2466 Prepared for Gisborne District Council, 30p.
- Phillips C, Marden M. (2016). Storm-initiated debris flows and plantation forestry: protocols for monitoring and post-storm data capture. Prepared for Gisborne District Council, 24p. Landcare Research Contract Report LC2607.
- Spiekermann, R & Marden, M. (2018). Best options for land use following radiata harvest in the Gisborne District under climate change: Spatial analysis of erosion susceptibility in plantation forests, East Coast Region SLMACC 405415. MPI Technical Paper No: 2018/47. ISBN No: 978-1-77665-953-1 (online) ISSN No: 2253-3923 (online).

## 9. Slash management and riparian buffers

- Baillie, BR., and Rolando, CA. (2015). Long-term management of streams in planted forest steeplands. *NZ Journal of Forestry*, Vol. 60, No. 2.
- Basher, L.R., Boothroyd, I.K.G., Quinn, J.M., Langer, E.R., Costley, K.J. and Steward, G. (2004). Riparian Buffers Mitigate Effects of Pine Plantation Logging on New Zealand Streams: 1. Riparian Vegetation Structure, Stream Geomorphology and Periphyton. *Forest Ecology and Management*, 194: 199–213.
- Quinn, J.M., Boothroyd, I.K.G. and Smith, B.J. (2004). Riparian Buffers Mitigate Effects of Pine Plantation Logging on New Zealand Streams: 2. Invertebrate Communities.



*Forest Ecology and Management*, 191: 129–146.

Rowe, D.K., Smith, J., Quinn, J. and Boothroyd, I. (2002). Effects of Logging With and Without Riparian Strips on Fish Species, Abundance, Mean Size, and the Structure of Native Fish Assemblages in Coromandel, New Zealand, Streams. *New Zealand Journal of Marine and Freshwater Research*, 36: 67–79.

## **10. Changes in mean bed levels East Coast rivers**

Hayward, K. A. (2001): Braided channel floodplain characteristics and development on the Tapuaeroa River, East Cape, New Zealand. MSc (Geography), University of Auckland. Pp.115. Key words: braided, channel, meandering, floodplain.

Peacock D. H., Marden M. 2004. Bed level changes in the Raparapaririki, Mangapoi and Mangawhairiki streams, Ruatoria. Engineering and Works Technical Report 2004/01 (EWTR 2004/01). 19p + 6 Appendices.

Marden M. 2005. Report to Gisborne District Council on Sediment Sources in Raparapaririki, Mangapoi and Mangawhairiki Streams, East Coast Region. Landcare Research Contract Report LC0304/088, pp. 10.

Peacock, D.H & Marden, M. (2013). Bed level changes in the Raparapaririki, Mangapoi and Mangawhairiki streams; Ruatoria volume 2: 2003 to 2013.

Peacock, D.H & Marden, M. (2019). Mean bed level trends in the upper Waipaoa River channel and Te Weraroa stream in response to land use change: 1948 to 2019. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

Peacock, D.H & Marden, M (2017). Mangatu River mean bed level trends; 1956 to 2017. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

Peacock, D.H (2017). Waiapu River mean bed level trends; 1958 to 2017. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council

Peacock, DH. (2017). Poroporo River mean bed level trends; 1958 to 2016. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

Peacock, D.H & Marden, M (2017). Mean bed level trends in the Manutahi and Mangaharei streams; Ruatoria 1979 to 2015. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

Peacock, D H (2017). Mangaoporo River mean bed level trends; 1958 to 2015. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

Peacock, D H (2017). Makatote-Kopuaroa Stream mean bed level trends;1958 to 2017. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

Peacock, D H (2017). Makarika Stream mean bed level trends; 1958 to 2017. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

Peacock, D H (2017). Paoaruku Stream mean bed level trends; 1958 to 2016. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

Peacock, D.H & Marden, M (2017). Karakatuwhero River mean bed level trends; 1968 to 2015. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

Peacock, D.H & Marden, M (2017). Whangaparaoa and Waikura rivers and the Mohau stream mean bed level trends; 1963 to 2019. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

Peacock, D.H & Marden, M (2017). Owēka Stream mean bed level trends; 1963 to 2019. Commissioned by the Environmental section and the (former) Roding section of the Gisborne District Council.

## 11. Sediment sources and delivery to stream channels

Allsop, F., 1973, *The Story of Mangatu: The Forest Which Healed the Land*. Wellington, New Zealand, A.R. Shearer, Government Printer, 100 p.

Betts, H.D., and De Rose, R.C., 1999, Digital elevation models as a tool for monitoring and measuring gully erosion: *International Journal of Applied Earth Observation and Geoinformation*, v. 1, no. 2, p. 91-101, [https:// doi .org /10 .1016 /S0303 -2434 \(99\)85002 -8](https://doi.org/10.1016/S0303-2434(99)85002-8).

Betts, H.D., Trustrum, N.A., and De Rose, R.C., 2003. Geomorphic changes in a complex gully system measured from sequential digital elevation models, and implications for management. *Earth Surface Processes and Landforms*, v. 28, no. 10, p. 1043-1058, [https:// doi .org /10 .1002 /esp .500](https://doi.org/10.1002/esp.500) .

Bilderback, E.L., Pettinga, J.R., Litchfield, N.J., Quigley, M., Marden, M., Roering, J.J., and Palmer, A.S., 2015, Hillslope response to climate-modulated river incision in the Waipaoa catchment, East Coast North Island, New Zealand. *Geological Society of America Bulletin*, v. 127, no. 1-2, p. 131-148. [https:// doi .org /10 .1130 /B31015 .1](https://doi.org/10.1130/B31015.1).

Chappell, P.R., 2016, *The Climate and Weather of the Gisborne District* (2nd ed.). National Institute of Water and Atmospheric Research Science and Technology Series Number 70, 40 p.

Derose, R., Gomez, B., Marden, M., and Trustrum, N., 1998. Gully erosion in Mangatu Forest, New Zealand, estimated from digital elevation models. *Earth Surface Processes and Landforms*, v. 23, no. 11, p. 1045-1053.

Fuller, I.C., and Marden, M., 2010, Rapid channel response to variability in sediment supply:

- Cutting and filling of the Tarndale fan, Waipaoa catchment, New Zealand. *Marine Geology*, v. 270, no. 1-4, p. 45-54.  
[https:// doi .org /10 .1016 /j.margeo .2009 .10 .004](https://doi.org/10.1016/j.margeo.2009.10.004)
- Fuller, I.C., and Marden, M., 2011. Slope-channel coupling in steepland terrain: A field-based conceptual model from the Tarndale gully and fan, Waipaoa catchment, New Zealand. *Geomorphology*, v. 128, no. 3-4, p. 105-115.  
[https:// doi .org /10 .1016 /j.geomorph .2010 .12 .018](https://doi.org/10.1016/j.geomorph.2010.12.018).
- Glade, T., 2003, Landslide occurrence as a response to land use change: A review of evidence from New Zealand. *Catena*, v. 51, no. 3-4, p. 297-314.  
[https://doi.org/10.1016/S0341-8162\(02\)00170-4](https://doi.org/10.1016/S0341-8162(02)00170-4)
- Kasai, M., Brierley, G.J., Page, M.J., Marutani, T., and Trustrum, N.A., 2005. Impacts of land use change on patterns of sediment flux in Weraamaia catchment, New Zealand: *Catena*, v. 64, no. 1, p. 27-60, [https:// doi.org /10 .1016 /j. Catena .2005 .06 .014](https://doi.org/10.1016/j.Catena.2005.06.014).
- Kasai, M. (2006). Channel processes following land use changes in a degrading steep headwater stream in North Island, New Zealand. *Geomorphology* 81 (3-4): 421-439.
- Leenman, A and Tunnicliffe, J. (2018). Genesis of a major gully mass-wasting complex, and implications for valley filling, East Cape, New Zealand. *GSA Bulletin*; v. 130; no. 7/8; p. 1121-1130.  
[https:// doi .org /10 .1130 /B31849 .1](https://doi.org/10.1130/B31849.1); 6 figures; 2 tables.
- Liebault, F., Gomez, B., Page, M., Marden, M., Peacock, D., Richard, D., and Trotter, C.M., 2005. Land-use change, sediment production and channel response in upland Regions. *River Research and Applications*, v. 21, no. 7, p. 739-756.  
[https:// doi .org /10 .1002 /rra.880](https://doi.org/10.1002/rra.880).
- Marden, M. (2005). Report to Gisborne District Council on Sediment Sources in Raparapaririki, Mangapoi and Mangawhairiki Streams, East Coast Region. Landcare Research Contract Report: LC0304/088.
- Marden, M.; Rowan, D. (1993): Protective value of vegetation on Tertiary terrain before and during Cyclone Bola, East Coast, North Island, New Zealand. *New Zealand Journal of Forestry Science* 23 (3): 255-263.
- Marden, M.; Rowan, D. (1999): Pre- and post-harvest site stability: Manutahi forest. Landcare Research Contract Report: LC9900/010.
- Marden, M., Arnold, G., Gomez, B., and Rowan, D., 2005. Pre- and post-reforestation gully development in Mangatu Forest, East Coast, North Island, New Zealand. *River Research and Applications*, v. 21, no. 7, p. 757-771.  
[https:// doi .org /10 .1002 /rra .882](https://doi.org/10.1002/rra.882).
- Marden, M., Herzig, A., and Arnold, G., 2011. Gully degradation, stabilisation, and effectiveness of reforestation in reducing gully-derived sediment, East Coast region, North Island, New Zealand. *Journal of Hydrology (New Zealand)*, v. 50, no. 1, p. 19-36.
- Marden, M., Arnold, G., Seymour, A., and Hambling, R., 2012. History and distribution of

- steepland gullies in response to land use change, East Coast region, North Island, New Zealand. *Geomorphology*, v. 153-154, p. 81-90.  
<https://doi.org/10.1016/j.geomorph.2012.02.011>.
- Marden, M., Herzig, A., and Basher, L., 2014, Erosion process contribution to sediment yield before and after the establishment of exotic forest: Waipaoa catchment, New Zealand. *Geomorphology*, v. 226, p. 162-174.  
<https://doi.org/10.1016/j.geomorph.2014.08.007>.
- Mazengarb, C.; Wilson, G. J.; Scott, G. H. (1991): A Miocene debris flow deposit, Puketoro Station, Raukumara Peninsula. New Zealand Geological Survey record 43: 107-111. Wellington. Department of Scientific & Industrial Research.
- Milliman, J.D.; Syvitski, P.M. (1992): Geomorphic/tectonic control of sediment discharge to the ocean: the importance of small mountainous rivers. *Journal of Geology* 100: 525-544.
- Miyamoto, A.; Marutani, T.; Kasai, M. (2003): Particle size distribution and sediment mobility in branch canals of the Waiapu Catchment, New Zealand. Spring Meeting of the Union, May 2003. Transactions, Japanese Geomorphological Union 24 (3). 324p.
- Marutani, T., Kasai, M., Reid, L.M., and Trustrum, N.A., 1999, Influence of storm-related sediment storage on the sediment delivery from tributary catchments in the upper Waipaoa River, New Zealand: *Earth Surface Processes and Landforms*, v. 24, no. 10, p. 881-896.  
[https://doi.org/10.1002/\(SICI\)1096-9837\(199909\)24:10<881::AID-ESP17>3.0.CO;2-I](https://doi.org/10.1002/(SICI)1096-9837(199909)24:10<881::AID-ESP17>3.0.CO;2-I).
- O'Byrne, T.N. (1967): A correlation of rock types with soils, topography and erosion in the Gisborne-East Cape region. *New Zealand Journal of Geology and Geophysics* 10: 217-231.
- Orpin, A. R.; Carter, L.; Kuehl, S. A.; Trustrum, N.A.; Lewis, K.B.; Alexander, C. R.; Gomez, B. (2002): Deposition from very high sediment yield New Zealand rivers is captured in upper margin basins. *Margins Newsletter* 9: 1-4. [www.nsf-margins.org/Publications/Newsletters/Issue9.pdf](http://www.nsf-margins.org/Publications/Newsletters/Issue9.pdf)
- Ota, Y.; Yoshikawa, T.; Moriya, I.; Iso, N.; Ikeda, Y.; Hull, A. G. (1985): River terraces of the eastern part of Raukumara Range, North Island, New Zealand. Unpublished manuscript. Copy held by the Institute of Geological & Nuclear Sciences Ltd.
- Page, M., Reid, L.M., and Lynn, I.H., (1999). Sediment production from Cyclone Bola landslides, Waipaoa catchment. *Journal of Hydrology (New Zealand)*, v. 38, no. 2, p. 289-308.
- Page MJ, Lukovic B 2011. An inventory of deep-seated landslides in the Waipaoa and Waimata catchments. GNS Science Report 2011/08, June 2011, 75p.
- Page M.J.; Trustrum N.A. (1997): A late Holocene lake sediment record of the erosion response to land use change in a steepland catchment, New Zealand. *Zeitschrift für*

Geomorphologie N.F. 41 (3): 369–392.

- Page, M., Marden, M., Kasai, M., Gomez, B., Peacock, D., Betts, H., Parkner, T., Marutani, T., and Trustrum, N. (2007). Changes in basin-scale sediment supply and transfer in a rapidly transformed New Zealand landscape. *Developments in Earth Surface Processes*, v. 11, p. 337–356, [https:// doi .org /10 .1016 /S0928 -2025 \(07\)11132 -9](https://doi.org/10.1016/S0928-2025(07)11132-9).
- Page, M.; Trustrum, N.; Gomez, B. (2000): Implications of a century of anthropogenic erosion for future land use in the Gisborne-East Coast Region of New Zealand. *New Zealand Geographer* 56 (2): 13-24.
- Page M.R.; Harmsworth G.R.; Trustrum N.; Kasai, M.; Muratani T. (2001): Waiapu River (North Island, New Zealand). In: Marutani, T.; Brierley, G.J.; Trustrum, N.A.; Page, M. (eds.) (2001): Source-to-sink sedimentary cascades in Pacific Rim Geo-Systems. Matsumoto Sabo Work Office, Ministry of Land, Infrastructure, and Transport. Motomachi, Matsumoto, Nagano, Japan, pp. 102–111.
- Page M.R.; Trustrum N.; Brackley H.; Gomez B.; Kasai M.; Muratani T. (2001): Waipaoa River (North Island, New Zealand). In: Marutani, T.; Brierley, G.J.; Trustrum, N.A.; Page, M. eds Source-to-sink sedimentary cascades in Pacific Rim geo-systems. Japan, Matsumoto Sabo Work Office, Ministry of Land, Infrastructure and Transport, Japan. Pp. 86–100.
- Parkner, T., Page, M.J., Marutani, T., and Trustrum, N.A. (2006). Development and controlling factors of gullies and gully complexes, East Coast, New Zealand. *Earth Surface Processes and Landforms*, v. 31, no. 2, p. 187–199. [https:// doi .org /10 .1002 /esp .1321](https://doi.org/10.1002/esp.1321).
- Parkner, T., Page, M., Marden, M., and Marutani, T. (2007). Gully systems under undisturbed indigenous forest, East Coast region, New Zealand. *Geomorphology*, v. 84, no. 3-4, p. 241–253. [https:// doi .org /10 .1016 /j.geomorph .2006 .01 .042](https://doi.org/10.1016/j.geomorph.2006.01.042).
- Pearce, A. J.; Black, R. D.; Nelson, C. S. (1981): Lithologic and weathering influences on slope form and process, eastern Raukumara Range, New Zealand. Pp 95-112. In: Davies, T. R. H. & Pearce, A. J. (eds.): Erosion and sediment transport in Pacific Rim steeplands. Washington. IAHS-AISH publication 132.
- Pearce, A. J. (1982): Complex mass-movement terrain in the Eastern Raukumara Peninsula, New Zealand: lithologic and structural-tectonic influences and some effects of recent deforestation and reforestation. In: Landslides and Mudflows and the Prevention of their Negative Impact on the Environment. (Proc. UNESCO Int. Seminar, Alma-Ata) Pp. 235-249.
- Pearce A. J.; O'Loughlin, C. L.; Jackson, R. J.; Zhang, X. B. (1987): Reforestation: on-site effects on hydrology and erosion, eastern Raukumara Range, New Zealand. In 'IAHS Publication no 167'.
- Phillips, C.J. (1988): Geomorphic effects of two storms on the upper Waitahaia River catchment, Raukumara Peninsula, New Zealand. *Journal of Hydrology (NZ)* 27 (2): 99-112.

- Phillips, C.J. (1989): Geomorphic effects of Cyclone Bola 1988 - A note. *Journal of Hydrology (NZ)* 28 (2): 142-146.
- Pick, M. C. (1962): The stratigraphy, structure and economic geology of the Cretaceous-Tertiary rocks of the Waiapu district, New Zealand. Unpublished Ph.D. thesis, University of Bristol, United Kingdom. Todd Oil Exploration Co.
- Rait, G. J. (1992): Early Miocene thrust tectonics on Raukumara Peninsula, northeastern New Zealand. Unpublished Ph.D. thesis, Victoria University of Wellington, Wellington, New Zealand.
- Rau, C. (1993): 100 years of Waiapu. Published by the Gisborne District Council, Gisborne Herald Co Ltd. 196 pp.
- Regnier, C. E.; Courtney, S. P.; Wiessing, M. I. (1988): Pukeamaru Ecological District. Survey Report for the Protected Natural Areas Programme No. 8. Department of Conservation, Wellington, New Zealand. 104p.
- Reid, L.M., and Page, M.J. (2002), Magnitude and frequency of landsliding in a large New Zealand catchment: *Geomorphology*, v. 49, no. 1, p. 71-88.
- Rijkse, W.C. (1980): Soils and agriculture of Waiapu Valley, East Coast, North Island, New Zealand. New Zealand Survey Report 60. 86 p. Maps 1.
- Rosser B 2007. Bank erosion in the Waikohu River: results of a field survey and possible causes of instability. Landcare Research Contract Report LC0708/085.
- Smith, R. K. (1974): Earthflows in the Poverty Bay-East Coast Region. Report on project NA/HY/5. Hydrological Research: Progress Report No. 18. Published by the Ministry of Works and Development for the National Water and Soil Conservation Organisation, New Zealand. 20p.
- Taylor, R.J., Massey, C., Fuller, I.C., Marden, M., and Archibald, G., (2017). Quantifying sediment connectivity in an actively eroding gully-mass movement complex, Waipaoa catchment, New Zealand. *Geomorphology*, <https://doi.org/10.1016/j.geomorph.2017.10.007>.
- Tunncliffe, J., Brierley, G., Fuller, I., Leenman, A., Marden, M., and Peacock, D. (2017). Reaction and relaxation in a coarse-grained fluvial system following catchment wide disturbance: *Geomorphology*, <https://doi.org/10.1016/j.geomorph.2017.11.006>.
- Trustrum, N.A.; Blaschke, P.M.; DeRose, R.C.; West, A. (1990): Regolith changes and pastoral productivity declines following deforestation in steeplands of North Island, New Zealand. Proceedings 14th International Congress of Soil Science, Kyoto, Japan. Pp. 125-130.
- Wadman, H. M.; McNinch, J. E.; Kuehl, S. A. (2006): Fine sediment sequestration on an active inner shelf, Waiapu River, New Zealand. American Geophysical Union, Fall Meeting 2006.
- Winkler, G. E. (1994): Engineering geological characterisation of Cretaceous - Late Tertiary rock units and their influence of landsliding, Waiapu District, East Cape. Thesis:

Master of Science. University of Canterbury.

- Wright, L.D.; Ma, Y.; Scully, M.; Friedrichs, C. T. (2005): Observations of storm driven benthic flows on the continental shelf off the Waiapu River, New Zealand. New Zealand Marine Sciences Society Meeting, Wellington, New Zealand, 23-26 August.
- Wright, L.D.; Ma, Y.; Scully, M.; Friedrichs, C. T. (2006): Observations of across shelf sediment transport during high energy flood events off the mouth of the Waiapu River, New Zealand. AGU 2006 Ocean Sciences Meeting, Honolulu, Hawaii, 20-24 February.
- Wright, L.D.; Friedrichs, C. T. (2006): Gravity-driven sediment transport on continental shelves: A status report. *Continental Shelf Research* 26 (17-18): 2092-2107.
- Yanxia, M. L.; Wright, D.; Friedrichs, C. T. (2007): Observations of sediment transport on the continental shelf off the mouth of the Waiapu River, New Zealand: Evidence for current-supported gravity flows. *Continental Shelf Research* (In Press, Accepted Manuscript).
- Yoshikawa, T.; Ikeda, Y.; Iso, N.; Moriya, I.; Hull, A.G.; Ota, Y. (1988): Origin and age of erosion surfaces in the upper drainage basin of Waiapu River, northeastern North Island, New Zealand. *New Zealand Journal of Geology and Geophysics* 31: 101-109.
- Zhang, X.; Phillips, C.J.; Marden, M. (1991): Internal deformation of a fast-moving earthflow, Raukumara Peninsula, New Zealand. *Geomorphology* 4: 145-154.
- Zhang, X.; Phillips, C.J.; Pearce, A. (1991): Surface movement in an earthflow complex, Raukumara Peninsula, New Zealand. *Geomorphology* 4 (3-4): 261-272.
- Zhang, X; Phillips, CJ; Marden, M. (1993). A comparison of earthflow movement rates on forested and grassed slopes, Raukumara Peninsula, North Island, New Zealand. *Geomorphology*, 6, 175- 187. [https://doi.org/10.1016/0169-555X\(93\)90045-4](https://doi.org/10.1016/0169-555X(93)90045-4)

## 12. Social implications of erosion and forestry

- Awatere S, Harmsworth G. (2012). Māori values –Iwi/Hapū perspectives of freshwater management in the Auckland region. Hamilton, Landcare Research.
- Awatere S, Harmsworth G, Pauling C. (2013). Using mātauranga Māori to inform freshwater planning. Lincoln, Landcare Research.
- Awatere, S., Marden, M., Warmenhoven, T., Pohatu, Daigneault, A., Monge, J., Dowling, L., Harrison, D (2018). Climate Resilient Māori Land. Contract Report: LC3133.
- Aldwell, P. H. B. (1982). Impacts of large scale forestry on settlement patterns in Waiapu County. *Planning Quarterly* 68, pp. 6-14.
- Aldwell, P. H. B. (1984). Some social and economic implications of large-scale forestry in Waiapu County. FRI Bulletin No.75. Forest Research Institute, Rotorua.



Fairweather & colleagues (Auckland University) have produced several papers on forestry-related impacts, including physical, economic, and social, of past Government policies on East Coast communities.

Harmsworth, G.R. 1995: Maori values for land use planning. Discussion document. Unpublished Manaaki Whenua-Landcare Research Report. 118 p.

Harmsworth, G.R. 1997: Maori values for land-use planning. Broadsheet newsletter of New Zealand Association of Resource Management, February 1997:37–52.

Harmsworth G.R. 1998: Indigenous values and GIS: a method and framework. *Indigenous Knowledge and Development Monitor* 6 (3): 3–7.

Harmsworth, G.R. 2001: A collaborative research model for working with iwi: discussion paper for FRST. Landcare Research Contract Report LC 2001/119. 29 p.

Harmsworth G, Raynor B. (2006). Cultural consideration in landslide risk perception. In: Glade T, Anderson MG, Crozier MJ ed. *Landslide hazard and risk*. Chichester, UK, John Wiley & Sons.

Harmsworth G, Tipa G 2009. Māori environmental monitoring in New Zealand: Progress, concepts, and future direction: Report for the ICM website. Palmerston North, Landcare Research. 29 p.

Harmsworth G, Awatere S. (2013). Indigenous Māori knowledge and perspectives of ecosystems. In: Dymond J, Ausseil A-G eds *Ecosystem services: conditions and trends*. Christchurch, Manaaki Whenua Press.

Harmsworth G, Warmenhoven T, Pohatu P, Page M. (2002). Waiapu catchment Technical Report: Maori community goals for enhancing ecosystem health. Landcare Research Contract Report LC0102/100 for Te Whare Wananga o Ngāti Porou, Ruatoria (Foundation for Research, Science, and Technology contract TWWX0001) (unpublished).

Warmenhoven, T. (2002): Waiapu research knowledge tables. Ruatorea, TWWONP.

### **13. Research posters**

Giadrossich, F., Marden, M., Urru1, M., Marrosu, R., Schwarz, M., Phillips, C., Cohen, D., Niedda, M. (2017). Bio-engineering traits of *Pinus radiata* D.Don. Presented at European Geosciences Union Assembly, Vienna, Austria.

Reidler, RA., Fuller, IC., Glade, T., Bell, R., Marden, M. (2013). Landslide driven erosion: Quantifying slope-channel coupling in forested steep terrain from 1946-2011, Tamaki catchment, New Zealand. Abstract for European Geosciences Union General Assembly, Vienna, Austria, 10-12 April 2013.

Taylor, R., Massey C., Fuller IC, Marden M, McSaveney M, Archibald G, Palmer N. (2012). Quantification of process in an actively eroding gully-mass movement complex, Waipaoa catchment, New Zealand. Poster EP31C-0828. American Geophysical Union Meeting, San Francisco, 3-7 December 2012.

Palmer A, Marden M, Litchfield N. (2011). Tephra from source to sink on an active plate margin: a chronology for events in the Waipaoa Catchment, East Coast North Island, New Zealand. Presented at INQUA Conference Bern, Switzerland 2011.

Marden, M., & Phillips, C. (c.2011). Growth performance of seral indigenous plant species during the first 5-years after establishment-plant trial, East Coast region, New Zealand.



Kahikatea\_final.pdf



Kauri\_final.pdf



Matai\_final.pdf



Miro\_final.pdf



Puriri\_final.pdf



Rimu\_final.pdf



Titoki\_final.pdf



Totara\_final.pdf

Fuller, I., & Marden, M. (2009). ‘Slope-channel coupling at a critical nexus in the Waipaoa sediment cascade: Tarndale Gully and Fan, Mangatu Forest, New Zealand’. Towards Integration and Synthesis of MARGINS S2S Research in PNG and NZ Focus Areas, 5-9 April 2009, Gisborne, New Zealand.

Holt, K., Marden, M, Palmer, A. (2009). ‘Vegetation history in the Gisborne –Waipaoa region and the relationship to post-glacial downcutting: a preliminary investigation from two sites in the Waimata River valley’. Towards Integration and Synthesis of MARGINS S2S Research in PNG and NZ Focus Areas, 5-9 April 2009, Gisborne, New Zealand.

Bender, C; Marsaglia, C; Vazquez, J; Mazengarb, C; & Marden, M. (2009). ‘A Mafic Conglomerate of Tectonic Significance, North Island, New Zealand’. Towards Integration and Synthesis of MARGINS S2S Research in PNG and NZ Focus Areas, 5-9 April 2009, Gisborne, New Zealand.

Carrasco, J; Marsaglia, K; Marden, M; Kirby, M. (2009). ‘Correlation of Physical, Mineralogical and Organic Properties of Late-Pleistocene Lacustrine Deposits from Redpath Paleolake, North Island, New Zealand’. Towards Integration and Synthesis of MARGINS S2S Research in PNG and NZ Focus Areas, 5-9 April 2009, Gisborne, New Zealand.

Marsaglia, K; James, D; De Vaughn, A; Parra, J; Rivera, K; Marden, M; & Kuehl, S. (2009). ‘Sediment Provenance Studies in the Waipaoa River Sedimentary System’. Towards Integration and Synthesis of MARGINS S2S Research in PNG and NZ Focus Areas, 5-9 April 2009, Gisborne, New Zealand.

Fuller, I., Marden, M., Massey, C. (2009). Connectivity in steepland environments: complex gully-fan interactions in the Tarndale system, Waipaoa catchment, New Zealand. 7th International Conference on Geomorphology ANZAIG); Ancient Landscapes-Modern Perspectives, 6-11 July, Melbourne, Australia.

Harmsworth, G.R.; Pohatu, P. (2000): Maori community goals for enhancing ecosystem health. Project poster. Te Whare Wananga o Ngati Porou and Manaaki Whenua.

Harmsworth, G.R.; Warmenhoven, T. (2001): "**Waiapu** Koka Huhua: Waiapu mother of many" Project poster. Te Whare Wananga o Ngati Porou and Manaaki Whenua. (Poster)

Harmsworth, G.; Warmenhoven, T. (2001): Ko **Waiapu** Te Awa art competition 2001 poster. Te Whare Wananga o Ngati Porou and Manaaki Whenua. (Poster)

Marden, M., Rowan, D., Phillips, C. (c.2006). Growth performance of common indigenous riparian plant species during the first 5-years after establishment-plant trial, East Coast region, New Zealand.



Riparianposter\_fina  
l\_Cabbage Tree.pdf



Riparianposter\_fina  
l\_Fivefinger.pdf



Riparianposter\_fina  
l\_Karamu.pdf



Riparianposter\_fina  
l\_Kohuhu.pdf



Riparianposter\_fina  
l\_Kowhai.pdf



Riparianposter\_fina  
l\_Lacebark.pdf



Riparianposter\_fina  
l\_Lemonwood.pdf



Riparianposter\_fina  
l\_Mapou.pdf



Riparianposter\_fina  
l\_Rewarewa.pdf



Riparianposter\_fina  
l\_Ribbonwood.pdf



Riparianposter\_fina  
l\_Tutu.pdf

Marden M, Phillips C. 2013. Survival and growth of poplar and willow pole plantings on East Coast hill country: a pilot study. LC 1622 Prepared for Plant & Food Research 16 pp.

Watson, A., & Marden, M. (2004). Root tensile strength as an indicator of performance of indigenous riparian plants-How do they work? Prepared for ICM AGM, December 2004, Nelson.

Marden, M., Rowan, D., Phillips, C. (2004). Performance of native riparian plants-how different are they? Prepared for ICM AGM, January 2004.

Marden, M., Rowan, D., Phillips, C. (2003). Performance of native riparian plants-how different are they? Prepared for ICM AGM, January 2003.

Page, M., Trustrum, N., Marden, M., Gomez, B. (2002). Erosion and sedimentation response to Holocene climate change, East Coast, North Island, New Zealand. American Geophysical Union conference, Wellington, June 2002.

Berryman, K.; Marden, M.; Mazengarb, C.; Rowan, D. (2000). Terraces and alluvial deposits of the Waipaoa and Waimata catchments: What have we learnt from studying them? Presented at Eco2000 Conference, Gisborne.

Phillips, C.; Fransen, P.; Marden, M.; Rowan, D. (2000). Sediment generation following harvesting of *Pinus radiata* in New Zealand: a variable story.

Phillips, C.; Watson, A.; Ekanayake, J.; Marden, M.; Rowan, D. (c. 1990). Are trees effective for erosion control? How do they work?

Phillips C & Marden M. 2013. Future forest erosion species trial-final report. LC 1626 Prepared for Future Forests Research Ltd, 19 pp.

Phillips, C.; Marden, M.; Rowan, D.; Ekanayake, J. (c. 1990). Riparian vegetation and erosion control: Value and function.

Rowe, L.; Marden, M.; Rowan, D. (c. 1990). Kanuka for erosion control; Why is it successful? The water story.

Marden, M.; Rowan, D.; Ekanayake, J.; Watson, A. (c. 1990). Kanuka for erosion control; Why is it successful? The root story.

Marden, M.; & Rowan, D. (c. 1990). Hauler Logging: the dirty story.

## 14. Popular articles

Basher, L.R. Phillips, C., Marden, M. and Harrison, D. (2015). What Do We Need for a Risk Management Approach to Steepland Plantation Forests in Erodible Terrain? *New Zealand Journal of Forestry*, 60(2): 7–10.

Davies-Colley, R.; Scarsbrook, M.; Marden, M. (1999): What lives in muddy east coast streams? Research in soft-rock catchments. *Water & Atmosphere* 7 (3): 7.

Davies-Colley, R.; Scarsbrook, M.; Marden, M. (1999). What lives in muddy streams? *Tairāwhiti Conservation Quorum*, Issue 18, p.4.

Freeman, T.; Marden, M., et al. (1999). Exploding the myth of the Tarndale Slip. *Conservation Quorum*, Issue 17. P.1-7.

Marden, M. (2003): Gully erosion, the cancer of Waiapu catchment: 901 reasons to act now. *Tairāwhiti Conservation Quorum*, Issue 32, pgs 1, 5-6, Winter 2003.

Marden, M, Basher, L., Phillips, C. and Black, R. (2015). Should Detailed Terrain Stability or Erosion Susceptibility Mapping Be Mandatory in Erodible Steep Lands? *New Zealand Journal of Forestry*, 59: 32–42.

Marden M & Phillips C. (2010). Growth performance of selected New Zealand indigenous conifer and broadleaved trees. *Tairāwhiti Conservation Quorum*, Issue No 61, 9-11.

Marden, M. (2006). Sediment sources following harvesting. *Conservation Quorum*, 43 Autumn 2006.

Marden, M & Phillips, C. (2006). Native Riparian Plant Research: an update. *Conservation Quorum*, 43, 6-7.

Marden, M. (2006). Expert says every tree adds value and protection. *Country-Wide*.

Marden, M. (2009). Gullies supply most sediment to major east coast river systems. *New Zealand Tree Grower*, 28-30, August 2009.

Marden, M., & Phillips, C. (2009). Native Plant Trial No. 2: how quickly do our native trees, grasses and herbs grow? *Tairāwhiti Conservation Quorum* 54, 6-8.

- Marden, M., & Phillips, C. (2009). Native riparian plants for riverbank stabilisation. *Catchment Connections* 5, 12-14, June 2009.
- Marden, M. (2007). Core- it' another paleo-lake! *Soil Horizons* 16, pg 4, September 2007.
- Marden, M. (2004). (article credited to Louise Thomas) Mapping ancient lands and predicting tidal waves. *Position Magazine*, 54-55.
- Marden, M. (2004). Native riparian plant trials - the final chapter. *Tairawhiti Conservation Quorum*, Issue 36, 7-8.
- Marden, M.; Phillips, C.; Rowan, D. (2003): Riparian plant trial: which species were the best performers? *Tairawhiti Conservation Quorum*, Issue 32, 12-13, Winter 2003.
- Marden, M. (2003). Researchers measure growth of native youngsters. *Discovery*.
- Marden, M.; & Phillips, C. (2002). How do native plants grow? *Tairawhiti Conservation Quorum*, Issue 26, p.14-15.
- Marden, M.; & Phillips, C. (2000). Using native plants to provide stability to streambanks. *Tairawhiti Conservation Quorum* Issue 21, p.6.
- Marden, M. (1999). Waipaoa river terraces: How old are they? *Tairawhiti Conservation Quorum*, Issue, 18, p.3.
- Marden, M.; Rowan, D.; Ekanayake, J.; Watson, A. (1999). Kanuka for erosion control-it really works! *Tairawhiti Conservation Quorum*, Issue 16, p.11.
- Marden, M; & Saunders H. (1992). Harvest at Mangatu: terrain on a tightrope. *New Zealand Forest Industries*, August, 60-61.
- Marden, M; & Saunders H. (1992). Balancing forestry economics and stewardship of the East Coast environment. *Conservation Quorum* 8, 8-9.
- Marden, M. (1989). East Coast Forests-a special case. *Forestry Forum*, 4, 6-7.
- Payn, T., Phillips, C., Basher, L., Baillie, B., Garrett, L., Harrison, D., Heaphy, M., Marden, M. (2015). Improving management of post-harvest risks in steepland plantations. *NZ Journal of Forestry*, August 2015, Vol. 60, No. 2. 6p.
- Phillips, C., Marden, M. and Basher, L. (2012). Plantation Forest Harvesting and Landscape Response – What We Know and What We Need to Know. *New Zealand Journal of Forestry*, 56: 4–12.
- Phillips, CJ, & Marden M. (2012). Future forest species for erosion control - something different? *Conservation Quorum* May 2012. <http://www.gdc.govt.nz/assets/Files/CQuorum/CQ-Autumn-2012.pdf>
- Phillips C, & Marden M. (2011). Soil stabilising characteristics of native plants. *Indigena*, November 2011: 10-13.

Phillips, C.; Ekanayake, J.; Marden, M. (1999). Can planting pattern make a difference to slope stability? *Tairawhiti Conservation Quorum*, Issue 17, p.14.

Phillips, C & Marden, M. (2006). Soil Conservation-is the renaissance coming? *Soil Horizons* 14, September 2006, pg. 2.

Scott, N.; Townsend, J.; Marden, M. (1999). Could scrub be worth dollars as a carbon sink? *Tairawhiti Conservation Quorum*, Issue 18, p.10-11.

Warmenhoven, T. (2000): "River research reveals serious sediment pollution". Te Runanga O Ngati Porou newsletter, NatiLink, October 2000.

## **15. Newspaper articles/Radio/TV**

Marden, M. (1998): New focus favoured for gully erosion. *Gisborne Herald*, 6 January 1998.

Marden, M. (1998). Area's erosion may shock world. *Gisborne Herald, Focus on the Land*, December 30.

Marden & Curtis. Something's got to change. *The Gisborne Herald* 19/ 01/ 2023.

Continuing scourge of gully erosion. *Gisborne Herald* 04 August 2020.

Slip, sliding, away. Michael Neilson. *Gisborne Herald Weekender*. 08/July 2017.

End of a Research era. Michael Neilson. 28 June 2017

Land use issues his focus. Profile on Dr Mike Marden. *Gisborne Herald*. 2015.

Presentation on gully erosion as part of 'Rural Delivery' programme on TV One, Saturday May 26th, 2012, 9 am.

Article on poplar research in *Gisborne Herald* 'Natures Land Anchor', Saturday 26 May 2012.

Interview highlighting gully erosion problems and solutions using East Coast research as an example. Screened on TV One 'Rural Delivery' on 26 May 2012.

Marden, M. (2007). Future-proof gullies researcher advises. *Gisborne Herald-Focus on the Land*, Wednesday 10 January 2007.

Marden, M. (2007). Core samples reveal the existence of an old lake. *Gisborne Herald*, 14th April 2007.

Marden, M. (2007) Drilling down to core the Waipaoas turbulent past. *Gisborne Herald*. January 10, 2007.

Marden, M. (2007). Erosion rate likely to increase-planting eroded land can earn carbon credits. *Gisborne Herald*, 27 June 2007.

Marden, M. (2004). Tarndale Slip highlights value of forestry. *Gisborne Herald* 24 April 2004.

Marden, M. (1999). New focus favoured for gully erosion. *Gisborne Herald*, January 6<sup>th</sup>.

Marden, M (1990). Some farms left unproductive. *The Hawkes Bay Herald Tribune*. November 30, 1990

Marden, M (1990). Erosion-the East Coast killer. *The Hawkes Bay Herald Tribune*. November 9, 1990.

Marden, M. (1998). Ash deposits are clue to history of Waipaoa River. *Gisborne Herald*, 11.05.98.

## 16. Soils

Basher L, McNeill S, Page M, Lynn I, Betts H, De Rose R, Marden M, Rosser B. 2013. Soil carbon stocks and changes: carbon losses from erosion. MPI Technical Paper no 2013/, 42 pp

Pullar WA, Penhale HR 1970. Periods of recent infilling of the Gisborne Plains basin. *New Zealand Journal of Science* 13: 410–434.

Rijkse, W.C. (1980): Soils and agriculture of Waiapu Valley, East Coast, North Island, New Zealand. *New Zealand Survey Report* 60. 86 p. Maps 1.

## 17. Geological maps

Blom, W. M. (1982): **Sedimentology of the Tokomaru Formation**, Waiapu Subdivision, Raukumara Peninsula. Thesis: Master of Science. University of Auckland.

Blom, W. M. (1984): **Stratigraphy and sedimentology** of Tokomaru Formation (Late Miocene - Early Pliocene), eastern Raukumara Peninsula, New Zealand. *New Zealand Journal of Geology and Geophysics* 27: 125-137.

Coleman, J. L. (1999): **Early Quaternary geology** of the Gisborne district (*Abstract*). Geological Society of New Zealand annual conference, Massey University. Geological Society of New Zealand miscellaneous publication 107A: 32.

Coleman, J. L. (1999): **Early-mid Quaternary geology** of the Gisborne district, East Coast, North Island, New Zealand. Unpublished M.Sc. thesis, University of Auckland, Auckland, New Zealand.

Driver, G. R. (1974): New Zealand **Land Resource Inventory** Gisborne – East Coast Region: **land use capability** extended legend. National Water and Soil Conservation Organisation, Wellington.

Eyles, G.O. (1985): The New Zealand **Land Resource Inventory erosion classification**. Water and Soil Miscellaneous Publication 85. 61 p.



- Gibbs, H. S. (1960): Soils of Gisborne – East Coast District and their problems for pastoral use. New Zealand DSIR, Soil Bureau Publication 215 (reprinted from N.Z. Grassland Proceedings).
- Harris, C. S.; McKee, J. G. (1964): Land Use capability survey, Gisborne/East Coast region, North Island, New Zealand. **Land use capability classes**: Sheets 1 and 2, scale one inch to two miles. N. Z. Department of Agriculture.
- Henderson, J.; Ongley, M. (1920): The Geology of the Gisborne and Whatatutu Subdivisions, Raukumara Division. Bulletin No. 21 (New Series), Geological Survey Branch, Department of Mines.
- Hewitt, A.E. 1998: New Zealand **Soil Classification**. Landcare Research Science Series 1, 2<sup>nd</sup> ed. Lincoln, Manaaki Whenua Press. 133 p.
- Jessen, M.R.; Crippen, T.; Page, M.J.; Rijkse, W.C.; Harmsworth, G.R.; McLeod, M. (1999): **Land Use Capability classification** of the Gisborne – East Coast region: A report to accompany the second-edition New Zealand Land Resource Inventory. Landcare Research Science Series No. 21. Lincoln, Manaaki Whenua Press.
- Kingma, J.T. (1964): Sheet 9 Gisborne, 1st edition. **Geological Map** of New Zealand, 1:250 000. Wellington, Department of Scientific and Industrial Research.
- Landcare Research (1999): New Zealand **Land Resource Inventory** at 1:50 000 for the Gisborne–East Coast Region, 2<sup>nd</sup> edition. Palmerston North, GIS database, Landcare Research.
- Mazengarb, C. (1993): **Cretaceous stratigraphy** of Raukumara Peninsula. Institute of Geological & Nuclear Sciences science report 93/20. Lower Hutt. Institute of Geological & Nuclear Sciences Ltd.
- Mazengarb, C.; Harris, D. H. M. (1994): **Cretaceous stratigraphic and structural relationships** of Raukumara Peninsula, New Zealand; stratigraphic patterns associated with the migration of a thrust system. *Annales Tectonicae* 8: 100-118.
- Mazengarb, C.; Speden, I.G. (compilers) (2000): **Geology** of the Raukumara area. IGNS 1:250 000 Geological Map 6. 1 Sheet + 60 p. Lower Hutt, Institute of Geological and Nuclear Sciences.
- Moore, P.R.; Francis, D.A.; Mazengarb, C. (1989): **Geological map of New Zealand** 1:250 000. DSIR Sheet QM 303 Raukumara, June 1989. New Zealand Geological Survey Report G138. Unpublished report and accompanying 1:250 000 map and a series of 1:50 000 field sheets.
- Moore, P.R.; Mazengarb, C. (1992): **Geology and landforms** of Raukumara Peninsula. In: Soons, J.M.; Selby, M.J. (eds.) *Landforms of New Zealand*, 2<sup>nd</sup> edn. Auckland, Longman Paul. Pp. 334–343.

New Zealand Forest Service (NZFS) (1971): FSMS6 Sheet No. 6 Raukumara, 1<sup>st</sup> edn. **Forest Class Map 1**: 250 000. Wellington, Forest Research Institute, New Zealand Forest Service.

Neef, G.; Bottrill, R. S. (1992): The **Cenozoic geology** of the Gisborne area (1:50 000 metric sheet Y18AB), North Island, New Zealand: with an appendix: clay minerals, quartz, plagioclase, and calcite in some sediments of the Tolaga Group. *New Zealand journal of geology and geophysics* 35:515-531.

Ongley, M.; Macpherson, E. O. (1928): The **Geology** of the **Waiapu Subdivision**, Raukumara Division. N.Z. Geol. Survey. Bull. 30. 79 pp.

Pullar, W.A. (1962): **Soils and Agriculture** of the Gisborne Plains. Soil Bureau Bulletin 20. New Zealand Department of Scientific and Industrial Research. Wellington, N.Z.: 91 pp.

## 18. **Waiapu Catchment, East Coast Region, North Island.**

*Keywords: rivers, vegetation, erosion, land use, forestry, storms, sediment*

- 1) Addington, L. D.; Kuehl S. A.; Mc Ninch, J. E. (2007): Contrasting modes of shelf sediment dispersal off a high-yield river: **Waiapu River**, New Zealand. *Marine Geology* 243 (1-4): 18-30.

Abstract: Recent studies of continental margins suggest that small, high-yield rivers are capable of generating shelf sediment-gravity flows, an idea that fundamentally alters our understanding of material flux from the continents to the ocean. Discharge measurements indicate that the Waiapu River, North Island, New Zealand reaches hyperpycnal concentrations ( $> 36 \text{ kg m}^{-2}$ ) on a yearly basis. This study contrasts shelf-edge basins with a broad trough along the shelf-edge off the Waiapu River, testing whether there is evidence that shelf sediment-gravity flows propagate to topographic lows. Observations and measurements through geochemical and sedimentological analyses of sediment cores, EM1002 swath bathymetry, and Chirp sub-bottom profiles suggest differing transport modalities on the outer shelf. In general, a southern trough-shaped region exhibits high terrigenous inputs and non-steady-state  $^{210}\text{Pb}$  profiles, whereas the northern basins contain steady-state  $^{210}\text{Pb}$  profiles and increased marine influence. Sediment-gravity flows dominate accumulation in the southern region, whereas within the northern portion, surface plume sedimentation is indicated. Overall, this study suggests that sediment-gravity flows could be bypassing the northern basins, perhaps a result of oceanographic influences and bathymetric steering as they seek a more direct route across the shelf.

Keywords: sedimentation; sediment-gravity flow; hyperpycnal flow; Waiapu River; New Zealand

- 2) Aldwell, P. H. B. (1982): Impacts of large-scale forestry on settlement patterns in Waiapu County. *Planning Quarterly* 68, pp. 6-14.
- 3) Aldwell, P. H. B. (1984): Some social and economic implications of large-scale forestry in **Waiapu County**. FRI Bulletin No.75. Forest Research Institute, Rotorua.

- 4) Betts, H.D.; DeRose, R.C. (1999): Digital elevation models as a tool for monitoring and measuring gully erosion. *Journal of Applied Earth Observation and Geoinformation*, 1(2): 91–101.
- 5) Betts, H.D.; Trustrum, N.A.; DeRose, R.C. (2003): Geomorphic changes in a complex gully system measured from sequential digital elevation models, and implications for management. *Earth Surface Processes and Landforms* 28 (10): 1043-1058.

Abstract: High-resolution digital elevation models (DEMs) were derived from sequential aerial photography of an active fluvio-mass movement (gully) complex in **New Zealand's North Island East Coast** region, to measure geomorphic changes over approximately one year. The gully showed a complex behaviour, combining fluvial and mass movement erosion, deposition, and reworking of materials stored in an active debris fan. During the measurement period  $5200 \pm 1700 \text{ m}^3$  of material were eroded from the 8.7 ha gully complex and  $670 \pm 180 \text{ m}^3$  from the 0.8 ha depositional fan, giving a total of  $5870 \pm 1710 \text{ m}^3$  for the entire gully complex-fan system.

The results provide a high-resolution description of gully behaviour over a short time period, and also demonstrate that mass movement (slumping and debris flows) accounted for almost 90 per cent ( $4660 \pm 200 \text{ m}^3$ ) of the sediment generated. This erosional response is described in terms of gully evolution by comparing the gully complex to other systems in the region in various stages of development. The effect of gully evolution on geomorphic coupling between the gully complex and channel system is described, and coupling is also shown to vary with the magnitude and frequency of rainfall events.

From a land management perspective the success of strategies, such as tree planting, to mitigate against gully erosion depends on the stage of gully development - particularly on whether or not mass movement erosion has begun. In contrast to gully rehabilitation efforts elsewhere, basin-wide afforestation in the early stages of gully incision is favoured over riparian planting, given that mass movement assisted by excessive groundwater pressure is the main process leading to uncontrollable gully expansion.

To protect land effectively against continuing gully erosion of headwater catchments and resulting downstream aggradation, it is necessary for land managers to understand the spatial and temporal variability of gully development fully so that mitigation efforts can be targeted appropriately.

- 6) Campbell, D. A. (1946): Down to the sea in slips. Soil Conservation and Rivers Control Council, Wellington, N. Z., Bulletin No. 5, 35p.
- 7) Campbell, D. A.; Campbell, A. P.; Hogg, R. J.; Scott, R. H.; Thomson, A. P.; Todd, A. D.; Vignaux, G. A.; Taylor, N. H. (1970): Wise land use and community development. Report of Technical Committee of Inquiry into the Problems of the Poverty Bay – East Cape District New Zealand. Published for the National Water and Soil Conservation Organisation by the Water and Soil Division, Ministry of Works Wellington, New Zealand. 120 p.
- 8) East Coast Project (1978): Report of land use planning and development study for erosion-prone land of the East Cape region, Section 1, The East Coast, May 1978. A report by the Poverty Bay Catchment Board, the 'Red Report'. 24p.
- 9) Eyles, G.O. (1983): The distribution and severity of present soil erosion in New Zealand. *New Zealand Geographer* 39 (1): 12–28.

- 10) Friedrichs, C. T.; Wright, L. D. (2004): Gravity-driven sediment transport on the continental shelf: implications for equilibrium profiles near river mouths. *Coastal Engineering* 51 (8-9): 795-811.  
 Abstract: An analytical model is developed for equilibrium bathymetric profiles off river mouths associated with the shoreward, convex upward portion of subaqueous deltas and clinoforms. The model builds on recent field results demonstrating that gravity-driven flux of suspended mud is important on shelves provided that wave-induced suspension of sediment supports the requisite turbid hyperpycnal layer. Because the maximum sediment load is determined by the critical Richardson number, the results are independent of the properties of the suspended mud or the bed. The model assumes the equilibrium state to represent a balance between the supply of sediment by a river at the coast and the downslope bypassing of sediment to deep water within wave-supported turbid near-bed layers. Progressive seaward increases in bed slope across the convex shelf profile allow the attenuation of wave agitation with depth to be compensated for by a downslope increase in the contribution of gravity. The model is consistent with shelf profiles off the mouths of the Eel (California), Ganges-Brahmaputra (Bangladesh), **Waiapu** (New Zealand), Po (Italy), and Rhone (France) Rivers. The equilibrium profile is predicted to be a function of wave climate and riverine sediment supply only, with deeper and broader profiles associated with decreasing sediment supply, increasing wave height and/or increasing wave period.  
 Keywords: Gravity-driven; Sediment transport; Continental shelf; Equilibrium profiles; Analytical model; River mouths
- 11) Gage, M.; Black, R. D. (1979): Slope Stability and geological investigations at **Mangatu State Forest**. Technical paper No. 66. Forest Research Institute, New Zealand Forest Service. 37pp. and maps.
- 12) Gibb, J. G. (1981): Coastal hazard mapping as a planning technique for **Waiapu County**, East Coast, North Island, New Zealand. He ripoata Whakature mo nga whenu papa-a-tai o te rohe o te kaunihera o Waiapu-Tairawhiti. Water & Soil technical publication. Wellington. Ministry of Works & Development.
- 13) Gibbs, H. S. (1954): Soils and Agriculture of Matakaoa County. New Zealand. Soil Bureau – Bulletin 11. New Zealand department of scientific and industrial research, 52p.
- 14) Gomez, B.; Eden, D.N.; Hicks, D.M.; Trustrum, N.A.; Peacock, D.H.; Wilmhurst, J. (1999): Contribution of floodplain sequestration to the sediment budget of the **Waipaoa river**, New Zealand. In: Marriott, S.B., Alexander, J. (eds) (1999): Floodplains: Interdisciplinary approaches. Special Publications, 163. London, Geological Society of London. Pp. 69–88.
- 15) Griffiths, G.A. (1982): Spatial and temporal variability in suspended sediment yields of **North Island basins**, New Zealand. *Water Resources Bulletin* 8 (4): 575-583.
- 16) Harmsworth, G.; Warmenhoven, T.; Pohatu, P. (2001): A **Waiapu catchment** management strategy: To stimulate interest and planning in the Waiapu, a strategy to underpin environmental management, economic development and sustain cultural values. A document for community and stakeholder discussion. Te Whare Wananga o Ngati Porou and Manaaki Whenua. 11p.

- 17) Harmsworth, G.; Warmenhoven T.; Pohatu, P.; Page, M. (2002): **Waiapu catchment** Technical Report: Maori community goals for enhancing ecosystem health. Foundation for Research, Science, and Technology (FRST) contract TWWX0001. Landcare Research report LC 0102/100 for Te Whare Wananga o Ngati Porou, Ruatorea (unpublished). 185p.
- 18) Hicks, D. M.; Griffiths, G. A. (1992): Sediment load. In: Mosley, M. P. (ed.): **Waters of New Zealand**. Wellington, New Zealand. Hydrological Society. Pp. 229–248.
- 19) Hicks, D.M.; Gomez, B.; Trustrum, N.A. (2000): Erosion thresholds and suspended sediment yields, Waipaoa River Basin, New Zealand. *Water Resources Research* 36 (4), 1129–1142.
- 20) Hicks, M.; Shankar, U.; McKerchar, A. (2003): Sediment yield estimates: a GIS tool. *Water & Atmosphere* 11(4): 26-27.  
<http://www.niwascience.co.nz/pubs/wa/ma/11-4/estimates>
- 21) Hicks, D.M.; Gomez, B.; Trustrum, N.A. (2004): Event suspended sediment characteristics and the generation of hyperpycnal plumes at river mouths: East Coast Continental Margin, North Island, New Zealand. *The Journal of Geology* 112: 471-485.  
 Abstract: Steepland rivers draining small, coastal watersheds often have very high suspended sediment loads, but the event characteristics of suspended sediment concentration and yield in this class of river is not well documented. Continuous monitoring at four sites in the Waipaoa River basin, New Zealand, demonstrates that during individual and composite events, suspended sediment concentration versus water discharge relations typically show clockwise hysteresis and that event maximum concentration is poorly related to event peak discharge. The signature of different erosion processes is also imprinted on the event yield magnitude frequency distributions. Gully-dominated tributary basins produce relatively high yields at all frequencies, reflecting greater sediment availability, whereas in tributary basins, where shallow landsliding is the dominant erosion process, there is a steep increase in yields in relation to return period. We estimate that flood discharges from the Waipaoa River approach or exceed the critical suspended sediment concentration ( $\sim 40,000$  mg L<sup>-1</sup>) for hyperpycnal plume generation (because of negative buoyancy) at the river mouth once every  $\sim 40$  yr, but in the neighbouring Waiapu and Uawa Rivers, the threshold concentration may be exceeded once a year and two to three times a year, respectively.
- 22) Hughes, S.; Hughes, I. (1990): The **Waiapu Mountains** of the Raukumara Range. Gisborne, Eastland Promotion Council/Te Rau Press.
- 23) Kasai, M.; Marutani, T.; Reid L.M.; Trustrum, N.A. (2001): Estimation of temporally averaged sediment delivery ratio using aggradational terraces in headwater catchments of the Waipaoa River, North Island, New Zealand. *Earth Surface Processes and Landforms* 26, 1–16.  
 Abstract: The sediment delivery ratio was estimated for two periods (28 years and eight years) following reforestation of seven tributary catchments (0.33 to 0.49 km<sup>2</sup>) in the headwaters of the Waipaoa River basin, North Island, New Zealand. In these catchments, gully erosion, which largely resulted from clearance of the natural forest between 1880 and 1920, is the main source of sediment to streams. Reforestation commenced in the early 1960s in an attempt to stabilize hillslopes and reduce sediment supply. Efforts have

been partially successful and channels are now degrading, though gully erosion continues to supply sediment at accelerated rates in parts of the catchment.

Data from the area indicate that the sediment delivery ratio (SDR) can be estimated as a function of two variables,  $\Psi$  (the product of catchment area and channel slope) and  $A_g$  (the temporally averaged gully area for the period). Sediment input from gullies was determined from a well defined relationship between sediment yield and gully area.

Sediment scoured from channels was estimated from dated terrace remnants and the current channel bed. Terrace remnants represent aggradation during major floods. This technique provides estimates of SDR averaged over periods between large magnitude terrace-forming events and with the present channel bed. The technique averages out short-term variability in sediment flux.

Comparison of gully area and sediment transport between two periods (1960-1988 and 1988-1996) indicates that the annual rate of sediment yield from gullies for the later period has decreased by 77 per cent, sediment scouring in channels has increased by 124 per cent, and sediment delivered from catchments has decreased by 78 per cent. However, average SDR for the tributaries was found to be not significantly different between these periods. This may reflect the small number of catchments examined. It is also due to the fact that the volume of sediment scoured from channels was very small relative to that produced by gullies.

According to the equation for SDR determined for the Waipaoa headwaters, SDR increases with increasing catchment area in the case where  $A_g$  and channel slope are fixed. This is because the amount of sediment produced from a channel by scouring increases with increasing catchment area. However, this relationship does not hold for the main stem of the study catchments, because sediment delivered from its tributaries still continues to accumulate in the channel. Higher order channels are, in effect, at a different stage in the aggradation/degradation cycle and it will take some time until a main channel reflects the effects of reforestation and its bed adjusts to net degradation.

Results demonstrate significant differences among even low order catchments, and such differences will need to be taken into consideration when using SDR to estimate sediment yields.

- 24) Kasai, M.; Brierley, G. J.; Page, M. J.; Marutani, T.; Trustrum, N. A. (2005): Impacts of land use change on patterns of sediment flux in **Weraamaia** catchment, New Zealand. *Catena* 64 (1): 27-60.

Abstract: Forest clearance between the 1890s and the early 1920s, subsequent scrub growth, and commencement of an afforestation program in 1979, modified the pattern and rate of sediment delivery to valley floors via shallow landslides and gully complexes in a steep headwater catchment (4.8 km<sup>2</sup>) in New Zealand. Analysis of the historical record, air photograph interpretation, and field survey indicates that both erosion types occurred across the catchment in the 1938 storm, aggrading channel beds and widening the active channel zone. In contrast, a 1 in 100 year event in 1988 (Cyclone Bola) induced numerous shallow landslides, but erosion of gully complexes was largely restricted to subcatchments that retained pasture, and the geomorphic impact of this event on channels was small. The changing volume and calibre of materials delivered to the valley floor, and the distribution of gully complexes, altered patterns and rates of channel adjustment after the events, and the resulting sediment flux. Development of gully complexes maintained coupling processes with channels for periods up to 10<sup>2</sup> years, forming wide channels in downstream reaches. Upstream–downstream connectivity along the trunk stream was altered by the formation of a large debris fan at the confluence with a tributary subjected to gully complex erosion. In contrast, slopes subjected to shallow landslides became decoupled from channels within 10 years, accelerating channel degradation and narrowing. Effective conveyance of a large volume of fine-grained materials promoted

immediate aggradation of gentle-gradient channels downstream. As gully complex areas stabilized following an increase in forest and scrub cover, channel courses became significant sediment sources. Although shallow landslide activity will continue to induce intermittent aggradation in the future, it is inferred that average sediment yield will continue to diminish to levels approaching those experienced prior to clearcutting, and the pattern of sediment flux will recover by 2030.

Keywords: Steep headwater catchment; Hillslope erosion; Storm events; Sediment budget; Coupling; Channel morphology; Land-use change

- 25) Kasai, M. (2006): Channel processes following land use changes in a **degrading steep headwater stream in North Island**, New Zealand. *Geomorphology* 81 (3-4): 421-439. Abstract: In headwater streams in steep land settings, narrow and steep valley floors provide closely coupled relationships between geomorphic components including hillslopes, tributary fans, and channel reaches. These relationships together with small catchment sizes result in episodic changes to the amount of stored sediment in channels. Major sediment inputs follow high magnitude events. Subsequent exponential losses via removal of material can be represented by a relaxation curve. The influence of hillslope and tributary processes on relaxation curves, and that of altered coupling relations between components, were investigated along a 1.3 km reach of a degrading channel in the 4.8 km<sup>2</sup> **Weraamaia Catchment**, New Zealand. Extensive deforestation in the late 19th and early 20th centuries, followed by invasion of scrubs and reforestation, induced changes to major erosion types from gully complexes to shallow landslides. Changes in the size and pattern of sediment slugs from 1938 to 2002 were analysed from air photographs tied to detailed field measurement. The rate and calibre of sediment flux changed progressively following substantive hillslope input in a storm in 1938. Subsequently, the channel narrowed and incised, decoupling tributary fans from the main stem, thereby scaling down the size of sediment slugs. As a consequence, the dominant influence on the behaviour of sediment slugs and associated relaxation processes, changed from tributary fans to the type and distribution of bedrock outcrops along the reach. Keywords: Channel morphology; Sediment slug; Relaxation process; Coupling processes; Land use change; Steep headwater catchments
- 26) Kenny, J. A. (1980): Geology of the Ihungia catchment, Raukumara Peninsula. M Sc, Dept. of Geology, University of Auckland.
- 27) Kniskern, T.A.; Kuehl, S. A. (2004): Sediment dispersal and deposition on the **Waiapu River Shelf**, N.Z., implications for sediment transport mechanisms and event preservation. American Geophysical Union Meeting, San Francisco, CA, December 2004.
- 28) Kniskern, T. A.; Harris, C. K.; Kuehl, S. A. (2006): Sediment deposition on the **Waiapu River Shelf**, N.Z., implications for sediment transport mechanisms and event Preservation. Ocean Sciences Meeting, Honolulu, Hawaii.
- 29) Kuehl, S.A.; Pratson, L.; Addington, L.; Gerald, L.; Gerber, T.; Kniskern, T.; Miller, A.; Liu, P.; Carter, L.; Orpin, A. (2006): Contrasting Shelf Sediment Dispersal off Small Mountainous rivers: The **Waipaoa and Waiapu Rivers**, NZ. Ocean Sciences Meeting, Honolulu, Hawaii.
- 30) Landcare Research (2001): **Waiapu project** GIS tables. Palmerston North, Landcare Research.



- 31) Leathwick, J. R.; Clarkson, B. D.; Burns, B. R. Innes, J. G. Smale, M. C. (1995): **Waiapu Ecological District**. Survey report for the Protected Natural Areas Programme. NZ PNA Programme No. 31. Department of Conservation, Gisborne.177 p.
- 32) Liébault, F.; Gomez, B.; Page, M.J.; Marden, M.; Peacock, D.H.; Richard, D.; Trotter, C.M. (2005): Land-use change, sediment production and channel response in upland regions. *River Research and Applications* 21, 739–756.
- 33) Marden, M.; Rowan, D. (1993): Protective value of vegetation on Tertiary terrain before and during Cyclone Bola, East Coast, North Island, New Zealand. *New Zealand Journal of Forestry Science* 23 (3): 255–263.
- 34) Marden, M.; Rowan, D. (1999): Pre- and post-harvest site stability: Manutahi forest. Landcare Research Contract Report: LC9900/010.
- 35) Marden, M. (2005): Report to Gisborne District Council on Sediment Sources in Raparapaririki, Mangapoi and Mangawhairiki Streams, East Coast Region. Landcare Research Contract Report: LC0304/088.
- 36) Marden, M.; Arnold, G.; Gomez, B.; Rowan, D. (2005): Pre- and post- reforestation gully development in Mangatu Forest, East Coast, North Island, New Zealand. *River Research and Applications* 21, 757–771.
- 37) Mazengarb, C.; Wilson, G. J.; Scott, G. H. (1991): A Miocene **debris flow** deposit, Puketoro Station, Raukumara Peninsula. *New Zealand Geological Survey record* 43: 107-111. Wellington. Department of Scientific & Industrial Research.
- 38) Milliman, J.D.; Syvitski, P.M. (1992): Geomorphic/tectonic control of sediment discharge to the ocean: the importance of small mountainous rivers. *Journal of Geology* 100: 525-544.
- 39) Miyamoto, A.; Marutani, T.; Kasai, M. (2003): Particle size distribution and sediment mobility in branch cannels of the **Waiapu Catchment**, New Zealand. Spring Meeting of the Union, May 2003. *Transactions, Japanese Geomorphological Union* 24 (3). 324p.
- 40) O’Byrne, T.N. (1967): A correlation of rock types with soils, topography and erosion in the Gisborne-East Cape region. *New Zealand Journal of Geology and Geophysics* 10: 217-231.
- 41) Orpin, A. R.; Carter, L.; Kuehl, S. A.; Trustrum, N.A.; Lewis, K.B.; Alexander, C. R.; Gomez, B. (2002): Deposition from very high sediment yield **New Zealand rivers** is captured in upper margin basins. *Margins Newsletter* 9: 1-4. [www.nsf-margins.org/Publications/Newsletters/Issue9.pdf](http://www.nsf-margins.org/Publications/Newsletters/Issue9.pdf)
- 42) Ota, Y.; Yoshikawa, T.; Moriya, I.; Iso, N.; Ikeda, Y.; Hull, A. G. (1985): River terraces of the eastern part of **Raukumara Range**, North Island, New Zealand. Unpublished manuscript. Copy held by the Institute of Geological & Nuclear Sciences Ltd.

- 43) Page M.J.; Trustrum N.A. (1997): A late Holocene lake sediment record of the erosion response to land use change in a **steep land catchment, New Zealand**. *Zeitschrift für Geomorphologie N.F.* 41 (3): 369–392.
- 44) Page, M. J.; Reid, L. M.; Lynn, I. H. (1999): Sediment production from Cyclone Bola landslides, **Waipaoa catchment**. *Journal of Hydrology (New Zealand)* 38(2): 289–308.
- 45) Page, M.; Trustrum, N.; Gomez, B. (2000): Implications of a century of anthropogenic erosion for future land use in the Gisborne-East Coast Region of New Zealand. *New Zealand Geographer* 56 (2): 13-24.  
 Abstract: The East Coast of the North Island of New Zealand is world renowned for its severe erosion, flooding, and sedimentation. Extensive deforestation between 1880-1920 initiated this period of dramatic landscape transformation, and today reforestation is seen as the panacea. However, a century of pastoral farming has left a legacy of a highly degraded landscape, which is currently redistributing the products of this erosion. The rate and level of landscape recovery will influence the ability of communities to carry out future land use. This paper uses the results of a decade of geomorphic research into the controls and processes of landscape change to illustrate some of the likely future impacts on the landscape and its land use, and to identify some still unanswered questions. This increasing understanding, together with changing community attitudes, provides the opportunity to maximise the benefits of reforestation and other management interventions.
- 46) Page M.R.; Harmsworth G.R.; Trustrum N.; Kasai, M.; Muratani T. (2001): **Waiapu River** (North Island, New Zealand). In: Marutani, T.; Brierley, G.J.; Trustrum, N.A.; Page, M. (eds.) (2001): *Source-to-sink sedimentary cascades in Pacific Rim Geo-Systems*. Matsumoto Sabo Work Office, Ministry of Land, Infrastructure, and Transport. Motomachi, Matsumoto, Nagano, Japan, pp. 102–111.
- 47) Page, M. J.; Marden, M.; Kasai, M.; Gomez, B.; Peacock, D. H. Betts, H. D.; Parkner, T.; Marutani, T.; Trustrum, N. A. (2007): Changes in basin-scale sediment supply and transfer in a rapidly transformed New Zealand landscape. In: Rinaldi, M.; Habersack, H.; Piegay, H. (eds.): *Developments in Earth Surface Processes 11, Gravel Bed Rivers 6: From Process Understanding to the Restoration of Mountain Rivers*, Chapter 13, 335-356, Elsevier B. V.
- 48) Parkner, T.; Page, M. J.; Marutani, T.; Trustrum, N. A. (2006): Development and controlling factors of gullies and gully complexes, East Coast, New Zealand. *Earth Surface Processes and Landforms* 31 (2): 187-199. Abstract: Sequential aerial photographs of a small headwater catchment in the **Waiapu** basin, East Coast Region, North Island, New Zealand, were interpreted to measure and analyse temporal changes in active area of gullies and gully complexes for a longer time span (1939-2003) and with higher temporal resolution compared to previous studies. We focus on the conditions leading to the development of gullies and gully complexes under pasture and forest by using topographic thresholds (slope-area relationships) of catchments for the initiation of gullies and gully complexes. In addition, the influence of two different lithologies as well as the occurrence of major rainfall events was related to gully activity. Twenty gullies and four gully complexes (occupying 62.5 ha or 12.5 per cent of the catchment area) occurred in the study catchment between 1939 and 2003. However, the majority of these were not active at all of the dates studied. Gullies developed in the

sandstone-dominated Tapuwaeroa Formation tended to attain their maximum size by 1957 with a mean catchment area of 2.1 ha. Gullies developed in mudstone of the Whangai Formation attained their maximum size in 1939 with a mean catchment area of 4.31 ha. Exceptions are gullies which developed into mass movement deposits or into an earth flow deposit as well as gullies developed under indigenous forest. Topographic threshold values for gullies under pasture and indigenous forest show that values for gullies under forest plot far above the threshold line of gullies under pasture, indicating that the topographical threshold for gully development under forest is higher compared to under pasture. A threshold value of 9.4 ha in catchment area is needed for the development of gully complexes under pasture, all located in the Whangai Formation and with the same orientation as the strike of the mudstones. Gully-complex area and dominance of mass-movement erosion increased with larger catchment area. A decreasing distance to the threshold line for gullies under pasture indicates a later development for gully complexes. No gully complexes developed under indigenous forest, indicating that the threshold value for gully-complex development is higher than for gully complexes under pasture and was not reached in the study area. A model of shifting topographical threshold for gully development for a given catchment is developed which depends on land use. When a catchment has an indigenous forest cover the topographical threshold is very high. After conversion to pasture, threshold values decrease drastically. With the invasion of scrub, the threshold slowly increases and returns to a similar level to that under indigenous forest after reforestation. Development of gullies and gully complexes is a highly dynamic phenomenon, and phases of expansion and inactivity indicate that models describing only unidirectional advancing stages without periods of inactivity are not suitable. Therefore, this study adds more phases to models of gully and gully-complex development in the East Coast Region. The threshold line for gully initiation under pasture and a value of 9.4 ha in catchment area for gully-complex initiation permits one to predict which catchments, under similar environmental settings, develop gullies and gully complexes on a physical basis. This enables land managers to implement sustainable land-use strategies to reduce erosion rates of gullies and gully complexes.

Keywords: gully erosion, gully complex, topographic threshold, land-use change

- 49) Parkner, T.; Page, M. J.; Marden, M.; Marutani, T. (2007): Gully systems under undisturbed indigenous forest, East Coast Region, New Zealand. *Geomorphology* 84: 241-253.
- 50) Parkyn, Davies-Colley, R.; Halliday, J.; Scarsbrook, M.; Nagels, J.; Marden, M.; Rowan, D. (2003): Environmental condition of streams under different land use on soft rock terrain. *New Zealand Journal of Marine and Freshwater Research* (??).
- 51) Parkyn, S. M.; Davies-Colley, R. J.; Halliday, N. J.; Scarsbrook, M. R.; Nagels, J. W., Marden, M.; Rowan, D. (2003). Environmental condition of streams in soft rock terrain. Does pine afforestation improve stream health in pasture streams? *NZ Journal of Marine and Freshwater Research* 32: 391- 605.
- 52) Pearce, A. J.; Black, R. D.; Nelson, C. S. (1981): Lithologic and weathering influences on slope form and process, **eastern Raukumara Range**, New Zealand. Pp 95-112. In: Davies, T. R. H. & Pearce, A. J. (eds.): *Erosion and sediment transport in Pacific Rim steeplands*. Washington. IAHS-AISH publication 132.
- 53) Pearce, A. J. (1982): **Complex mass-movement** terrain in the Eastern Raukumara Peninsula, New Zealand: lithologic and structural-tectonic influences and some effects of recent

deforestation and reforestation. In: Landslides and Mudflows and the Prevention of their Negative Impact on the Environment. (Proc. UNESCO Int. Seminar, Alma-Ata) Pp. 235-249.

- 54) Pearce A. J.; O'Loughlin, C. L.; Jackson, R. J.; Zhang, X. B. (1987): Reforestation: on-site effects on hydrology and erosion, eastern Raukumara Range, New Zealand. In 'IASH Publication no 167'.  
Keywords: earthflow/erosion/gully/hydrology/reforestation
- 55) Peacock, D. H.; Marden, M. (2004): Bed level changes in the Raparapaririki, Mangapoi, and Mangawhairiki Streams; Ruatoria. Engineering and Works Technical Report 2004/01. Gisborne District Council, Gisborne, New Zealand.
- 56) Phillips, C.J. (1988): Geomorphic effects of two **storms** on the upper Waitahaia River catchment, Raukumara Peninsula, New Zealand. Journal of Hydrology (NZ) 27 (2): 99-112.
- 57) Phillips, C.J. (1989): Geomorphic effects of **Cyclone Bola** 1988 - A note. Journal of Hydrology (NZ) 28 (2): 142-146.
- 58) Pick, M. C. (1962): The stratigraphy, structure and economic geology of the Cretaceous-Tertiary rocks of the **Waiapu district**, New Zealand. Unpublished Ph.D. thesis, University of Bristol, United Kingdom. Todd Oil Exploration Co. Ltd. Unpublished open-file petroleum report 608. Wellington. Ministry of Economic Development.
- 59) Rait, G. J. (1992): Early Miocene thrust tectonics on **Raukumara Peninsula**, northeastern New Zealand. Unpublished Ph.D. thesis, Victoria University of Wellington, Wellington, New Zealand.
- 60) Rau, C. (1993): 100 years of **Waiapu**. Published by the Gisborne District Council, Gisborne Herald Co Ltd. 196 pp.
- 61) Regnier, C. E.; Courtney, S. P.; Wiessing, M. I. (1988): Pukeamaru Ecological District. Survey Report for the Protected Natural Areas Programme No. 8. Department of Conservation, Wellington, New Zealand. 104p.
- 62) Richardson, J.; Jowett, I. G. (2002): Effects of sediment on fish communities in **East Cape streams**, North Island, New Zealand. New Zealand Journal of Marine and Freshwater Research (36): 431-442. <http://www.rsnz.org/publish/nzjmfr/2002/033.php>
- 63) Rosser B, Marden M, Harmsworth G. 2008: Literature review and catalogue of physical resource information for the Waiapu Catchment, East Coast Region, Landcare Research Contract Report: LC0708/118, prepared for Gisborne District Council and Foundation for Research Science and Technology, 109p.
- 64) Rijkse, W.C. (1980): Soils and agriculture of **Waiapu** Valley, East Coast, North Island, New Zealand. New Zealand Survey Report 60. 86 p. Maps 1.
- 65) Smith, R. K. (1974): Earthflows in the Poverty Bay-East Coast Region. Report on project NA/HY/5. Hydrological Research: Progress Report No. 18. Published by the Ministry of

Works and Development for the National Water and Soil Conservation Organisation, New Zealand. 20p.

- 66) Trustrum, N.A.; Blaschke, P.M.; DeRose, R.C.; West, A. (1990): Regolith changes and pastoral productivity declines following deforestation in steeplands of North Island, New Zealand. Proceedings 14<sup>th</sup> International Congress of Soil Science, Kyoto, Japan. Pp. 125–130.
- 67) Wadman, H. M.; McNinch, J. E.; Kuehl, S. A. (2006): Fine sediment sequestration on an active inner shelf, **Waiapu River**, New Zealand. American Geophysical Union, Fall Meeting 2006.
- 68) Walling, D.E.; Webb, B.W. (1996): Erosion and sediment yield: A global overview. IAHS Publication 236. Proceedings of the Exeter Symposium, July 1996, University of Exeter, England. Pp. 3–19.
- 69) Warmenhoven, T. (2002): **Waiapu** research knowledge tables. Ruatorea, TWWONP.
- 70) Winkler, G. E. (1994): Engineering geological characterisation of Cretaceous - Late Tertiary rock units and their influence of landsliding, **Waiapu District**, East Cape. Thesis: Master of Science. University of Canterbury.
- 71) Wright, L.D.; Ma, Y.; Scully, M.; Friedrichs, C. T. (2005): Observations of storm driven benthic flows on the continental shelf off the **Waiapu River**, New Zealand. New Zealand Marine Sciences Society Meeting, Wellington, New Zealand, 23-26 August.
- 72) Wright, L.D.; Ma, Y.; Scully, M.; Friedrichs, C. T. (2006): Observations of across shelf sediment transport during high energy flood events off the mouth of the **Waiapu River**, New Zealand. AGU 2006 Ocean Sciences Meeting, Honolulu, Hawaii, 20-24 February.
- 73) Wright, L.D.; Friedrichs, C. T. (2006): Gravity-driven sediment transport on continental shelves: A status report. *Continental Shelf Research* 26 (17-18): 2092-2107.  
 Abstract: Recent field observations from several shelf environments show that gravity-driven transport within negatively buoyant layers is an important mode of fine sediment transport across continental shelves. Specifically, Dick Sternberg, along with his students and colleagues, stimulated a paradigm shift by reporting strong evidence from the Amazon and Eel shelves that hyperpycnal layers do not require auto-suspension for sustenance but can be initiated by sediment flux convergence and supported by wave and current-induced suspension within relatively thin near-bed layers. As these layers move downslope under the influence of gravity, they may deposit sediment in response to decreases in bottom orbital velocities, near-bed current velocity, and/or bed slope. Direct or indirect evidence for wave or current supported sediment gravity flows has recently been reported off other high-load rivers including the Atchafalaya, Fly, Ganges–Brahmaputra, Klamath, Mad, Mississippi, Po, Rhone, **Waiapu**, **Waipaoa**, Yangtze, and Yellow among others. Growing evidence from observational and modelling studies suggests that flux convergence followed by wave and current supported gravity driven transport is a primary cause of across-shelf transport and emplacement of flood deposits on many muddy shelves and may be a major contributor to and control on the large-scale formation and morphology of subaqueous deltas and shelf clinofolds. Recent and

ongoing studies on this subject are synthesized in this paper and recommendations are offered for further study.

Keywords: Fluid mud; Sediment transport; Turbidity currents; Subaqueous delta

- 74) Yanxia, M. L.; Wright, D.; Friedrichs, C. T. (2007): Observations of sediment transport on the continental shelf off the mouth of the **Waiapu River**, New Zealand: Evidence for current-supported gravity flows. *Continental Shelf Research* (In Press, Accepted Manuscript).
- 75) Yoshikawa, T.; Ikeda, Y.; Iso, N.; Moriya, I.; Hull, A.G.; Ota, Y. (1988): Origin and age of erosion surfaces in the upper drainage basin of **Waiapu River**, northeastern North Island, New Zealand. *New Zealand Journal of Geology and Geophysics* 31: 101-109.
- 76) Zhang, X.; Phillips, C.J.; Marden, M. (1991): Internal deformation of a fast-moving earthflow, **Raukumara Peninsula**, New Zealand. *Geomorphology* 4: 145-154.  
 Abstract: Sub-surface deformation of a fast-moving earthflow was studied in the East Coast Region of the North Island, New Zealand, for a period of 2 years. Tiltmeter profiles indicated that earthflow materials were subjected to a variety of movement mechanisms, including internal deformation by gravity-shearing flow, extension flow, compression flow, rotation over curved slopes, rotation of material beneath the earthflow\*, and sliding. Overall, internal deformation accounted for less than 25% of the total surface movement, the remaining 75% being the result of sliding movement along the basal shear plane. Gravity-shearing flow occurs along the basal shear plane, which is thought not to be more than a few centimetres thick. Micro-topography (features on a scale of 1–10 m) largely affected tilting behaviour (internal deformation) of fastmoving earthflows. In the longitudinal direction, tiltmeter profiles on curved slopes were monoclinical and rotation was the principal mechanism causing tilt. On concave slopes the tiltmeter profile had a forward (downslope)-convex shape, with compression being the principal deformation mechanism. On even (planar) slopes, extension flow (creep) predominated, with the profile showing little evidence of rotation. However, minimal deformation (tilting) was likely to be caused by a combination of either sub-earthflow rotation, extension flow, or compression flow. In the lateral direction, topographic influences produced considerable deformation, particularly on concave slopes as a result of compression flow. There was little or no lateral deformation on even slopes.
- 77) Zhang, X.; Phillips, C.J.; Pearce, A. (1991): Surface movement in an earthflow complex, **Raukumara Peninsula**, New Zealand. *Geomorphology* 4 (3-4): 261-272.  
 Abstract: The surface movement of an unstable earthflow complex was studied over a 10-year period using a network of pegs inserted to a depth of 1 m. The mean and maximum surface movement velocities in the transport zones of the two study earthflows ranged from 0.2–0.4 m and 1.7–2.8 m/month respectively. Maximum velocities for individual pegs ranged up to 3 m/month. The movement data indicated that the earthflows were approximately in a steady state for the 10-year duration of the study. Application of Iverson's theory for the kinematics of unsteady, non-uniform landslide movement indicated that the earthflows had Pe values in the range of 0.1 to 2.0 (Pe is the dimensionless parameter known as the landslide Peclet number). These values indicated that the earthflows showed a blend of plastic and viscous behaviour and that diffusion might be more significant than kinematic-wave propagation in transmitting disturbances.

- 78) Zhang, X.; Phillips, C.J.; Marden, M. (1993): A comparison of earthflow movement rates on forested and grassed slopes, **Raukumara Peninsula**, North Island, New Zealand. *Geomorphology* 6: 175-187.

Abstract: Surface movement rates on forested earthflows are 2–3 orders of magnitude less than those on grassed earthflows. Subsurface deformation results largely from extension flow on grassed earthflows and from compression flow on forested earthflows. A rafting mechanism in which blocks of roots from individual trees interact with those of neighbouring trees to retard surface movement is used to explain deformation profiles of forested earthflows. The rheology of earthflow materials is changed by the presence of tree roots.

## 19. Waipaoa Catchment, North Island

Black RD 1977. Rivers of change: Early history of the Upper Waipaoa and Mangatu catchments. Christchurch, Protection Forestry Division, Forestry Research Institute.

Black, R. D. (1980): Upper Cretaceous and Tertiary geology of Mangatu State Forest, Raukumara Peninsula, New Zealand. *N. Z. Journal of Geology and Geophysics* 23: 293-312.

DeRose, R.C.; Gomez, B.; Marden, M.; Trustrum, N.A. (1998): Gully erosion in Mangatu Forest, New Zealand, estimated from digital elevation models. *Earth Surface Processes and Landforms* 23: 1045–1053.

Abstract: The methodology and errors involved in determining the amount of sediment produced during two (19.5 and 33.2 year) periods by 11 (c. 0.01 - >0.20 km<sup>2</sup>) gullies within a 4 km<sup>2</sup> area in the headwaters of the Waipaoa River basin, New Zealand, using sequential digital elevation models are described. Sediment production from all gullies within the study area was  $0.99 \pm 0.03 \times 10^6 \text{ t a}^{-1}$  ( $2480 \pm 80 \text{ t ha}^{-1} \text{ a}^{-1}$ ) during the period from 1939 to 1958. It declined to  $0.62 \pm 0.02 \times 10^6 \text{ t a}^{-1}$  ( $1550 \pm 50 \text{ t ha}^{-1} \text{ a}^{-1}$ ) during the period from 1958 to 1992, when many of the smaller gullies were stabilized by a programme of afforestation, which commenced in 1960. Both figures are very high by global standards. The two largest (the Tarndale and Mangatu) gully complexes together generated 73 and 95 per cent of the sediment in the specified time periods, but the latter amount is equivalent to only c. 5 per cent of the total annual sediment load of the Waipaoa River.

Trustrum, N.A. & Page, M.J. (2000). Paleoenvironmental signals of land use and climate change in the Lake Tutira and Waipaoa sedimentary systems, New Zealand. *Proceedings of the IGBP Land Use and Climate Change in Fluvial Systems, LUCIFS workshop, 20-21 March, 2000, Bonn, Germany.* (Abstract).

Gomez, B, Dennis N. Eden, D. Murray Hicks, Noel A. Trustrum, David H. Peacock & Janet Wilmshurst (1999). Contribution of floodplain sequestration to the sediment budget of the Waipaoa River, New Zealand. In: Marriott, S. B. & Alexander, J. (eds) 1999. *Floodplains: Interdisciplinary Approaches*. Geological Society, London, Special Publication 163, p 69-88.

Berryman, K, Michael Marden, Dennis Eden, Colin Mazengarb, Yoko Ota & Ichio Moriya (2000) Tectonic and paleoclimatic significance of Quaternary River terraces of the

Waipaoa River, East Coast, North Island, New Zealand. *New Zealand Journal of Geography and Geophysics* Vol. 43, p229-245.

- Eden, D.N, Alan S. Palmer, Shane J. Cronin, Michael Marden & Kelvin Berryman (2001). Dating the culmination of river aggradation at the end of the Last Glaciation using distal tephra compositions, eastern North Island, New Zealand. *Geomorphology* Vol. 32, p133-151.
- Carter, L., Manighetti, B., Elliot, M., Trustrum, N.A. & Gomez, B. Source, sea level, and circulation effects on the sediment flux to the deep ocean during marine isotope Stage 1 off eastern New Zealand. *Global and Planetary Change*.
- Gomez, B., Trustrum, N.A. & Hicks, D.M. (1998). Links between hillslope erosion and floodplain sequestration: Waipaoa River basin, New Zealand. In: *The American Geophysical Union, Fall Meeting, December 6 - 10, 1998, San Francisco, California.* p227. (Poster paper and Abstract).
- Trustrum, N.A., Gomez, B., Page, M.J., Reid, L.M., Hicks, D.M., Marutani, T., & Marden, M. (1998). Linking magnitude-frequency relations from hillslope to floodplain. In: *Eighth Biennial Conference of the Australian and New Zealand Geomorphology Group, Goolwa, South Australia, November 15-20, 1998, p57.* (Abstract).
- Marden, M., Berryman, K. (1999) Waipaoa River terraces: How old are they? *Conservation Quorum* 18, Spring 1999.
- Gomez, B. (2000) The MARGINS Source-to-Sink programme and its significance to the Gisborne East Coast area. *Eco2000 Workshop, Gisborne 7-8 March 2000.* (Abstract).
- Trustrum, N.A. (2000) The Waipaoa sedimentation project: regional relevance and global significance. *Eco2000 workshop, Gisborne 7-8 March 2000.* (Abstract).
- Page, M.J. & Trustrum, N.A. (2000) High resolution lake sediments from New Zealand - a record of Late Holocene storm history, vegetation change and landscape response. *PAGES Newsletter, Vol. 8, No. 3, December 2000.*
- Page, M, Noel Trustrum & Basil Gomez (2000) Implications of a century of Anthropogenic erosion for future land use in the Gisborne-East Coast region of New Zealand. *New Zealand Geographer* Vol. 56, No. 2, p9-20.
- Page M.R.; Trustrum N.; Brackley H.; Gomez B.; Kasai M.; Muratani T. (2001): Waipaoa River (North Island, New Zealand). In: Marutani, T.; Brierley, G.J.; Trustrum, N.A.; Page, M. eds *Source-to-sink sedimentary cascades in Pacific Rim geo-systems.* Japan, Matsumoto Sabo Work Office, Ministry of Land, Infrastructure and Transport, Japan. Pp. 86-100.
- Page, M, Noel Trustrum & Basil Gomez (2000). Legacy of a century of erosion for future land use in Eastland. *Eco2000 workshop, Gisborne 7-8 March 2000.* (Poster).
- Trustrum, N.A, B. Gomez, M. J. Page, L. M. Reid & D. M. Hicks (1999). Sediment production, storage and output: The relative role of large magnitude events in steepland catchments. *Zeitschrift fur Geomorphologie, Suppl.* Bd. 115, p71-86.



- Reid, L. M., & N.A. Trustrum (In Press) Sediment budgets and land management planning: examples from NZ. *Journal of Environmental Planning and Management*.
- Reid, L.M., & M.J. Page (In Press) Magnitude and frequency of landsliding in a large New Zealand catchment. *Geomorphology*.
- Gomez, B, Craig Fulthorpe, Lionel Carter, Kelvin Berryman, Greg Browne, Mal Green, Murray Hicks & Noel Trustrum (2001) MARGINS Source to Sink: New Zealand Focus Area. *EOS, Transactions, American Geophysical Union*. Vol 82, No 14, 3rd April 2001.
- Kasai, M, Tomomi Marutani, Leslie M. Reid & Noel A. Trustrum (2001) Estimation of temporally averaged sediment delivery ratio using aggradational terraces in headwater catchments of the Waipaoa River, North Island, NZ. *Earth Surface Processes and Landforms* Vol. 26, p1-16.
- Marden, M., Berryman, K., Eden, D., Mazengerb, C., Ota, Y. & Moriya, I. (2000) Tectonic and paleoclimatic significance of Quaternary river terraces of the Waipaoa River, East Coast, North Island, New Zealand. In: 9th Australia-New Zealand Geomorphology Group (ANZGG) Conference, Wanaka, Dec 2000, p5. (Abstract).
- Page, M.J., Trustrum, N.A. & Brackley, H.L. (2001) Erosion-related soil carbon losses in a steep-land catchment, New Zealand. 5th International Geomorphology Conference, August 2001, Tokyo.
- Fulthorpe, C., Gomez, B., Berryman, K., Carter, L., Hicks, M., and Trustrum, N. (2000). MARGINS and the Waipaoa Sedimentary System. Poster prepared for Eco2000 workshop, Gisborne, 7-8 March 2000. (Abstract).
- Hicks, D.M, Basil Gomez & Noel A. Trustrum (2000). Erosion thresholds and suspended sediment yields, Waipaoa River Basin, NZ. *Water Resources Research* Vol. 36, No. 4, p1129-1142.
- Page, M., Trustrum, N. & Brackley, H. (2000). Erosion-related soil carbon losses in the Tutira catchment since European settlement. In: 9th Australia-New Zealand Geomorphology Group (ANZGG) Conference, Wanaka, Dec 2000, p65. (Abstract).
- Tate, K.R, N.A. Scott, A. Parshotam, L. Brown, R.H. Wilde, D.J. Giltrap, N.A. Trustrum, B. Gomez & D.J. Ross (2000). A multi-scale analysis of a terrestrial carbon budget. Is New Zealand a source or sink of carbon? *Agriculture, Ecosystems and Environment* Vol. 82, p229-246.
- Trustrum, N. (2000). Controls on sediment generation in the Waipaoa Sedimentary System. Presentation at MARGINS Workshop, September 2000, Lake Tahoe, USA.
- Trustrum, N., Page, M., Gomez, B., Marden, M., Berryman, K. & Hicks, M. (2000). Landscape evolution and controls on sediment and soil carbon fluxes. In: 9th Australia-New Zealand Geomorphology Group (ANZGG) Conference, Wanaka, Dec 2000, p94. (Abstract).
- Trustrum, N., Brackley, H. & Page, M. (2000). Sediment budget research workshop and East Coast field trip. October 2000. Sediment budget workshop and field trip schedule for Japanese Delegation. East Coast, North Island, New Zealand.

- Page, M. (2000). Effectiveness of trees in reducing hill country erosion. MAF Resource Management Update Issue 4, April 2000, p7-8.
- Trustrum, N.A. and Page, M.J. (2000). Paleoclimate signals in the Lake Tutira and Waipaoa sedimentary systems (Abstract). Paleoclimate meeting organised by the NZ Royal Society, March 10, 2000. Wellington.
- Betts, H. D. & Ronald C. DeRose (1999). Digital Elevation Models as a tool for monitoring and measuring gully erosion. JAG Vol. 1, Issue 2, p91-101.
- Gomez, B., Trustrum, N.A., Hicks, D.M., & Page, M.J. (1999). Impact of anthropogenically-induced degradation on suspended sediment and carbon fluxes: the case of the Waipaoa River basin, New Zealand. Geological Society of America Annual Meeting, October 27-30, 1999, Denver, Colorado. (Abstract).
- Basil Gomez, Noel A. Trustrum & D. Murray Hicks (1999). Magnitude and frequency characteristics of suspended sediment transport and floodplain construction in the Waipaoa River Basin, New Zealand. Drainage Basin Dynamics Conference, Jerusalem, May 1999. (Abstract).
- Marutani, T.; Kasai, M.; Reid, L.M.; Trustrum, N.A. (1999): Influence of storm-related sediment storage on the sediment delivery from tributary catchments in the Upper Waipaoa River, New Zealand. *Earth Surface Processes and Landforms* 24: 881–896.
- Page, M.J., L.M. Reid & I.H. Lynn (1999). Sediment production from Cyclone Bola landslides, Waipaoa Catchment. *Journal of Hydrology* Vol. 38, No. 2, p289-308.
- Wilmshurst, J.M., Dennis N. Eden & Paul C. Froggatt (1999). Late Holocene forest disturbance in Gisborne, New Zealand: a comparison of terrestrial and marine pollen records. *New Zealand Journal of Botany* Vol. 37, p523-540.
- DeRose, R.C, Basil Gomez, Mike Marden & N. A. Trustrum (1998). Gully erosion in Mangatu Forest, New Zealand, estimated from Digital Elevation Models. *Earth Surface Processes and Landforms* Vol. 23, p1045-1053.

## **20. Waimata Catchment**